Title: *In vitro* and *in vivo* activity of hydrolyzed Saccharomyces cerevisiae against goat nematodes

Authors: Naylene C.S. Silva, Aldilene S. Lima, Carolina R. Silva, Danilo R.B. Brito, José A.A. Cutrim-Junior, Marlise N. Milhomem, Livio M. Costa-Junior

PII: S0304-4017(18)30084-0
DOI: https://doi.org/10.1016/j.vetpar.2018.02.034
Reference: VETPAR 8618

To appear in: *Veterinary Parasitology*

Received date: 13-12-2017
Revised date: 30-1-2018
Accepted date: 21-2-2018

Please cite this article as: Silva NCS, Lima AS, Silva CR, Brito DRB, Cutrim-Junior JAA, Milhomem MN, Costa-Junior LM, *In vitro* and *in vivo* activity of hydrolyzed Saccharomyces cerevisiae against goat nematodes, *Veterinary Parasitology* (2010), https://doi.org/10.1016/j.vetpar.2018.02.034

In vitro and *in vivo* activity of hydrolyzed *Saccharomyces cerevisiae* against goat nematodes

Naylene C. S. Silva[1], Aldilene S. Lima[2], Carolina R. Silva[3], Danilo R. B. Brito[4], José A. A. Cutrim-Junior[4], Marlise N. Milhomem[1], Livio M. Costa-Junior[1*]

[1]Universidade Federal do Maranhão, Programa de Pós-Graduação em Ciências da Saúde, Centro de Ciências Biológicas e da Saúde, São Luís, Maranhão, Brazil

[2]Programa de Pós-Graduação em Biotecnologia, RENORBIO, São Luís, Maranhão, Brazil.

[3]Programa de Pós-Graduação em Biodiversidade e Biotecnologia, BIONORTE, São Luís, Maranhão, Brazil.

[4]Instituto Federal do Maranhão, Curso de Zootecnia, São Luís, Maranhão, Brazil.

*Author correspondence

Tel.: +55 98 32729547

E-mail address: livio.martins@ufma.br; livioslz@yahoo.com

Highlights

- Hydrolyzed yeast cells are a byproduct of sugarcane alcohol production.
- The effect of hydrolyzed yeast on *H. contortus* was evaluated through *in vitro* and *in vivo* assays.
- The yeast inhibited larval exsheathment of the *H. contortus*.
- The yeast was able to reduce FECs in animals from D40 to D54, showing an average efficiency of 60%.