### **Accepted Manuscript**

Labrasol® and salts of medium chain fatty acids can be combined in low concentrations to increase the permeability of a macromolecule marker across isolated rat intestinal mucosae

Joanne Heade, Sam Maher, Sinead B. Bleiel, David J. Brayden

PII: S0022-3549(18)30096-0


DOI: 10.1016/j.xphs.2018.02.012

Reference: XPHS 1084

To appear in: Journal of Pharmaceutical Sciences

Received Date: 14 December 2017 Revised Date: 25 January 2018

Accepted Date: 9 February 2018



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



#### ACCEPTED MANUSCRIPT

# Labrasol® and salts of medium chain fatty acids can be combined in low concentrations to increase the permeability of a macromolecule marker across isolated rat intestinal mucosae

Joanne Heade<sup>1,3</sup>, Sam Maher<sup>2</sup>, Sinead B. Bleiel<sup>3</sup>, David J. Brayden<sup>1†</sup>

<sup>1</sup>UCD School of Veterinary Medicine and UCD Conway Institute, University College Dublin, Belfield, Dublin 4, <sup>2</sup>School of Pharmacy, Royal College of Surgeons in Ireland, St. Stephen's Green, Dublin 2, <sup>3</sup>AnaBio Technologies Ltd, IDA Business Park, Carrigtwohill, Co. Cork, Ireland.

†Corresponding author: Tel.: +3537166013, Email: david.brayden@ucd.ie

### Download English Version:

## https://daneshyari.com/en/article/8513280

Download Persian Version:

https://daneshyari.com/article/8513280

<u>Daneshyari.com</u>