Accepted Manuscript



Inhibition of mitochondrial permeability transition pore opening contributes to cannabinoid type 1 receptor agonist ACEA-induced neuroprotection

Lei Ma, Wen Niu, Shuai Yang, Junbin Tian, Hanlin Luan, Ming Cao, Wenbin Xi, Weifeng Tu, Ji Jia, Jianrui Lv

PII: S0028-3908(18)30135-7

DOI: 10.1016/j.neuropharm.2018.03.024

Reference: NP 7127

To appear in: *Neuropharmacology*

Received Date: 31 October 2017

Revised Date: 19 March 2018

Accepted Date: 20 March 2018

Please cite this article as: Ma, L., Niu, W., Yang, S., Tian, J., Luan, H., Cao, M., Xi, W., Tu, W., Jia, J., Lv, J., Inhibition of mitochondrial permeability transition pore opening contributes to cannabinoid type 1 receptor agonist ACEA-induced neuroprotection, *Neuropharmacology* (2018), doi: 10.1016/j.neuropharm.2018.03.024.

Inhibition of mitochondrial permeability transition pore opening contributes to cannabinoid type 1 receptor agonist ACEA-induced neuroprotection

Lei Ma[a,1], Wen Niu[b,1], Shuai Yang[c,1], Junbin Tian[a], Hanlin Luan[d], Ming Cao[d], Wenbin Xi[d], Weifeng Tu[d], Ji Jia[d,*], Jianrui Lv[a,*]

[1] These authors contributed equally to this work.

* Correspondence should be addressed to Jianrui Lv (Email: ljr2011@139.com, Tel: 86-29-87679599) and Ji Jia

(Email: jiaji1981@126.com, Tel: 86-20-88653504).

[a] Department of Anesthesiology, the Second Affiliated Hospital of Xi'an Jiaotong University, Xi'an, 710004, China

[b] Department of Pathology and Pathophysiology, the Fourth Military Medical University, Xi'an, 710032, China

[c] Department of Neurosurgery, Guangzhou General Hospital of Guangzhou Military Command, Guangzhou, 510010, China

[d] Department of Anesthesiology, Guangzhou General Hospital of Guangzhou Military Command, Guangzhou, 510010, China