

Accepted Manuscript

Activation of ERK/CREB pathway in noradrenergic neurons contributes to hypernociceptive phenotype in H4 receptor knockout mice after nerve injury

Maria Domenica Sanna, Tommaso Mello, Emanuela Masini, Nicoletta Galeotti



PII: S0028-3908(17)30492-6

DOI: [10.1016/j.neuropharm.2017.10.025](https://doi.org/10.1016/j.neuropharm.2017.10.025)

Reference: NP 6910

To appear in: *Neuropharmacology*

Received Date: 21 July 2017


Revised Date: 18 October 2017

Accepted Date: 20 October 2017

Please cite this article as: Sanna, M.D., Mello, T., Masini, E., Galeotti, N., Activation of ERK/CREB pathway in noradrenergic neurons contributes to hypernociceptive phenotype in H4 receptor knockout mice after nerve injury, *Neuropharmacology* (2017), doi: 10.1016/j.neuropharm.2017.10.025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Nerve injury



Download English Version:

<https://daneshyari.com/en/article/8517551>

Download Persian Version:

<https://daneshyari.com/article/8517551>

[Daneshyari.com](https://daneshyari.com)