Accepted Manuscript

Xanthone-rich extract from Gentiana dinarica transformed roots and its active component norswertianin induce autophagy and ROS-dependent differentiation of human glioblastoma cell line

Gordana Tovilovic-Kovacevic, Dijana Krstic-Milosevic, Branka Vinterhalter, Mina Toljic, Vladimir Perovic, Vladimir Trajkovic, Ljubica Harhaji-Trajkovic, Nevena Zogovic

 PII:
 S0944-7113(18)30087-4

 DOI:
 10.1016/j.phymed.2018.03.052

 Reference:
 PHYMED 52426

To appear in: *Phytomedicine*

Received date:3 November 2017Revised date:7 February 2018Accepted date:19 March 2018

Please cite this article as: Gordana Tovilovic-Kovacevic, Dijana Krstic-Milosevic, Branka Vinterhalter, Mina Toljic, Vladimir Perovic, Vladimir Trajkovic, Ljubica Harhaji-Trajkovic, Nevena Zogovic, Xanthone-rich extract from Gentiana dinarica transformed roots and its active component norswertianin induce autophagy and ROS-dependent differentiation of human glioblastoma cell line, *Phytomedicine* (2018), doi: 10.1016/j.phymed.2018.03.052

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Xanthone-rich extract from *Gentiana dinarica* transformed roots and its active component norswertianin induce autophagy and ROS-dependent differentiation of human glioblastoma cell line

Gordana Tovilovic-Kovacevic^a, Dijana Krstic-Milosevic^b, Branka Vinterhalter^b, Mina Toljic^c, Vladimir Perovic^d, Vladimir Trajkovic^d, Ljubica Harhaji-Trajkovic^{e,*}, Nevena Zogovic^{e,*}

^aDepartment of Biochemistry, Institute for Biological Research "Sinisa Stankovic", Despot Stefan Blvd 142, 11000 Belgrade, University of Belgrade, Serbia

^bDepartment of Plant Physiology, Institute for Biological Research "Sinisa Stankovic", Despot Stefan Blvd 142, 11000 Belgrade, University of Belgrade, Serbia

^cGenetic Laboratory Department, Obstetrics and Gynecology Clinic "Narodni Front", Street Kraljice Natalije 62, 11000 Belgrade, Serbia

^dInstitute for Microbiology and Immunology, School of Medicine, Dr Subotic Street 1, 11000 Belgrade, University of Belgrade, Serbia

^eDepartment of Neurophysiology, Institute for Biological Research "Sinisa Stankovic", Despot Stefan Blvd 142, 11000 Belgrade, University of Belgrade, Serbia

*Corresponding authors

Ljubica Harhaji-Trajkovic, Nevena Zogovic

Department of Neurophysiology, Institute for Biological Research "Sinisa Stankovic", Despot Stefan Blvd 142, 11000 Belgrade, University of Belgrade, Serbia

Tel.: +381 11 36 43 233; fax: +381 11 27 61 433

E-mail addresses: buajk@yahoo.com (Lj. Harhaji-Trajkovic); nevenar@ibiss.bg.ac.rs (N. Zogovic)

1

Download English Version:

https://daneshyari.com/en/article/8517928

Download Persian Version:

https://daneshyari.com/article/8517928

Daneshyari.com