Accepted Manuscript



Intradermal and intranasal immunizations with oligomeric middle layer rotavirus VP6 induce Th1, Th2 and Th17☐T cell subsets and CD4+T lymphocytes with cytotoxic potential

Suvi Heinimäki, Maria Malm, Timo Vesikari, Vesna Blazevic

PII: S0166-3542(18)30174-8

DOI: 10.1016/j.antiviral.2018.06.012

Reference: AVR 4316

To appear in: *Antiviral Research*

Received Date: 22 March 2018

Revised Date: 9 May 2018

Accepted Date: 19 June 2018

Please cite this article as: Heinimäki, S., Malm, M., Vesikari, T., Blazevic, V., Intradermal and intranasal immunizations with oligomeric middle layer rotavirus VP6 induce Th1, Th2 and Th17☐T cell subsets and CD4+ T lymphocytes with cytotoxic potential, *Antiviral Research* (2018), doi: 10.1016/j.antiviral.2018.06.012.

Intradermal and intranasal immunizations with oligomeric middle layer rotavirus VP6 induce Th1, Th2 and Th17 T cell subsets and $CD4^+$ T lymphocytes with cytotoxic potential

Suvi Heinimäki, Maria Malm, Timo Vesikari, Vesna Blazevic*

Vaccine Research Center, Faculty of Medicine and Life Sciences, University of Tampere, Finland

Email adresses:

suvi.heinimaki@uta.fi

maria.malm@uta.fi

timo.vesikari@uta.fi

vesna.blazevic@uta.fi

***Corresponding author**:

Vesna Blazevic, PhD

Biokatu 10

33520 Tampere

Finland

Email: vesna.blazevic@uta.fi