Accepted Manuscript



Inhibition of human cytomegalovirus replication by tricin is associated with depressed CCL2 expression

Yumiko Akai, Hidetaka Sadanari, Masaya Takemoto, Noboru Uchide, Tohru Daikoku, Naofumi Mukaida, Tsugiya Murayama

PII: S0166-3542(17)30541-7

DOI: 10.1016/j.antiviral.2017.09.018

Reference: AVR 4158

To appear in: *Antiviral Research*

Received Date: 28 July 2017

Revised Date: 8 September 2017

Accepted Date: 18 September 2017

Please cite this article as: Akai, Y., Sadanari, H., Takemoto, M., Uchide, N., Daikoku, T., Mukaida, N., Murayama, T., Inhibition of human cytomegalovirus replication by tricin is associated with depressed CCL2 expression, *Antiviral Research* (2017), doi: 10.1016/j.antiviral.2017.09.018.

Inhibition of human cytomegalovirus replication by tricin is associated with depressed CCL2 expression

Yumiko Akai[a], Hidetaka Sadanari[a], Masaya Takemoto[a], Noboru Uchide[b], Tohru Daikoku[a], Naofumi Mukaida[c], Tsugiya Murayama[a]*

[a]Department of Microbiology and Immunology, [b]Department of Clinical Pharmacy, Faculty of Pharmaceutical Sciences, Hokuriku University, Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
[c]Division of Molecular Bioregulation, Cancer Research Institute, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

* Corresponding author. Tel.: +81 76 229 6223; Fax: +81 76 229 2781.
E-mail address: t-murayama@hokuriku-u.ac.jp (T. Murayama).

Abbreviations: DMEM, Dulbecco's Modified Eagle's Medium; HCMV, human cytomegalovirus; HEL, human embryonic lung fibroblasts cells; IE, immediate early gene.