Accepted Manuscript



A 5-hydroxyoxindole derivative attenuates LPS-induced inflammatory responses by activating the p38-Nrf2 signaling axis

Tomomi Niino, Kenji Tago, Daisuke Yasuda, Kyoko Takahashi, Tadahiko Mashino, Hiroomi Tamura, Megumi Funakoshi-Tago

PII: S0006-2952(18)30234-X
DOI: https://doi.org/10.1016/j.bcp.2018.06.021
Reference: BCP 13175

To appear in: *Biochemical Pharmacology*

Received Date: 18 April 2018
Accepted Date: 14 June 2018

Please cite this article as: T. Niino, K. Tago, D. Yasuda, K. Takahashi, T. Mashino, H. Tamura, M. Funakoshi-Tago, A 5-hydroxyoxindole derivative attenuates LPS-induced inflammatory responses by activating the p38-Nrf2 signaling axis, *Biochemical Pharmacology* (2018), doi: https://doi.org/10.1016/j.bcp.2018.06.021

A 5-hydroxyoxindole derivative attenuates LPS-induced inflammatory responses by activating the p38-Nrf2 signaling axis.

Tomomi Niino[a], Kenji Tago[b], Daisuke Yasuda[c], Kyoko Takahashi[c], Tadahiko Mashino[c], Hiroomi Tamura[a], and Megumi Funakoshi-Tago[a]

[a]Department of Hygienic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan

[b]Division of Structural Biochemistry, Department of Biochemistry, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke-shi, Tochigi-ken 329-0498, Japan

[c]Division of Medicinal Chemistry and Bio-organic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan

*Corresponding author: Megumi Funakoshi-Tago

Tel.: +81-3-5400-2689, Fax: +81-3-5400-2689, E-mail: tago-mg@pha.keio.ac.jp

Department of Hygienic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan