



Tanshinone IIA inhibits cervix carcinoma stem cells migration and invasion via inhibiting YAP transcriptional activity

Jinghao Qin^a, Hongbing Shi^c, Yanjie Xu^c, Fang Zhao^c, Qing Wang^{b,*}

^a Traditional Chinese Medicine Department, Changzhou First People's Hospital, 185th on Juqian street, Changzhou, 213000, China

^b Department of Gynecology, Changzhou First People's Hospital, 185th on Juqian street, Changzhou, 213000, China

^c Department of Oncology, Changzhou First People's Hospital, 185th on Juqian street, Changzhou, 213000, China

ARTICLE INFO

Keywords:

HuR
YAP
Cervix carcinoma
Stemness
Adriamycin

ABSTRACT

This study aims to explore the effects and related mechanisms of Tanshinone IIA in cervix carcinoma (CC) stemness-like cells migration, invasion, stemness and chemotherapeutic sensitivity. Here, we found that Tanshinone IIA suppressed CC stemness-like cells migration and invasion in a concentration- and time-dependent manner. And consistent results were obtained in CC cells stemness characterized as the decrease of CC stemness markers expression and cells spheroid formation ability. Mechanistically, we found that Tanshinone IIA suppressed RNA binding protein HuR translocation from nuclear to cytoplasm, and thus reduced YAP mRNAs stability and transcriptional activity. Importantly, overexpression YAP-5SA rescued the inhibition of Tanshinone IIA on CC cells stemness. Furthermore, Tanshinone IIA enhanced adriamycin sensitivity in CC stemness-like cells, this effect was attenuated by YAP-5SA overexpression too. Therefore, Tanshinone IIA could suppress CC stemness-like cells migration and invasion by inhibiting YAP transcriptional activity.

1. Introduction

Cervical cancer (CC) is one of the common gynecological malignant tumors that seriously threaten women health, and is closely related with the human papillomavirus (HPV) infection and the HPV vaccine has achieved remarkable results in the prevention of cervical cancer [1]. Although surgery, radiotherapy and chemotherapy have exerted better effects on early CC treatment, the efficacy on local stage and metastatic CC is limited and the survival rate of recurrent CC is lower [2]. Cancer stem cells (CSCs) have been confirmed to be involved in tumor initiation, metastasis and recurrence [3], thus, finding novel drugs that could eliminate CC CSCs or suppress CC cells stemness may provide potential methods or new clues for treating CC patients.

Danshen (*Salvia miltiorrhiza Bunge*) has been used extensively and historically in China to treat various diseases, including cardiovascular diseases, cerebrovascular diseases and cancer [4]. TanshinoneIIA is one of the major monomer extracted from root of *Salvia miltiorrhiza* (RSM), which holds various activities, especially in tumors progression [5] and inflammation [6]. Previous studies have shown that TanshinoneIIA could inhibit the growth of glioma and breast cancer stem cells [7,8]. However, the roles and related mechanisms of Tanshinone IIA in CC stemness have never been reported.

Hippo pathway is a conservative signaling in mammals, and consists

of MST1/2 (mammalian Sterile 20-like kinase 1/2), LATS1/2 (large tumor suppressor 1/2) which could phosphorylate and inactivate the downstream transcriptional effectors YAP/TAZ [9]. YAP/TAZ has been shown to contribute to tumor cells stemness and inhibition of YAP/TAZ transcriptional activity could attenuate CSCs progression [10,11]. Notably, LATS1/2 could suppress breast cancer EMT and metastasis via inactivating YAP/TAZ activity [12]. However, effectors in facilitating the development of YAP/TAZ are frustrated. Therefore, it is an urgent need to find novel YAP/TAZ inhibitors.

RNA binding proteins are a kind of proteins that could bind to and enhance mRNAs stability [13]. HuR, as an RNA binding protein, has been shown to facilitate CSCs progression, such as HuR facilitates cancer stemness of lung cancer cells via regulating miR-873/CDK3 and miR-125a-3p/CDK3 axis [14], and miR-146b-5p overexpression attenuates stemness and radioresistance of glioma stem cells by targeting HuR/lincRNA-p21/beta-catenin pathway [15]. Here, we found that Tanshinone IIA could inhibit HuR translocation from nuclear to cytoplasm. Furthermore, we indicated that HuR directly bound to YAP and enhanced YAP transcriptional activity. Importantly, we showed that Tanshinone IIA attenuated CC cells stemness, CC stem cells migration and invasion in a concentration- and time- dependent manner, these effects were attenuated by YAP-5SA overexpression which could not be phosphorylated by LATS1/2. Finally, we found that Tanshinone IIA

* Corresponding author.

E-mail address: qing_wang@sohu.com (Q. Wang).

enhanced adriamycin sensitivity in CC cells. Therefore, our results suggest that Tanshinone IIA could suppress CC stem cells formation, migration and invasion by activating Hippo pathway.

2. Material and methods

2.1. Cells culture and reagents

CC cells Hela, and C33 A, and healthy primary normal cervical epithelial cells HcerEpic were purchased from the Chinese Academy of Sciences Cell Bank. All of the cell lines were cultured in Dulbecco's Minimum Essential Medium (DMEM) medium (Gibco, USA) with 10% FBS (fetal bovine serum, Gibco) plus 80 U/ml penicillin and 0.08 mg/ml streptomycin at 37 °C under humidified atmosphere with 5% CO₂. Tanshinone IIA (Cat # S2365) and adriamycin (S1208) were purchased from Selleck.cn. pQCXIH-Myc-YAP-5SA (YAP-5SA) plasmid (Plasmid # 33093), 8xGTIIc-luciferase plasmid (Plasmid # 34615), a YAP-responsive synthetic promoter driving luciferase expression plasmid and pFRT_TO_eGFP_ELAVL1 (Plasmid #106105), a plasmid inducing HuR expression, were purchased from Addgene.

2.2. Transfection

Cells were seeded at the density of 5×10^5 cells/well in a 6-well plate. After 24 h, 2.5 µg of YAP-5SA plasmid was transfected into cells using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) following the manufacturer's instructions before Tanshinone IIA treatment.

2.3. Cell spheroid formation assay

CC cells with YAP-5SA transfection or not were cultured in ultra-low attachment 24-well plates (Corning, Union City, CA) at 500 cells/well with Mammocult™ Human Medium Kit (Cat#05620, Stemcell Technologies, Vancouver, BC, Canada), followed by Tanshinone IIA treatment. After 10 days culture, mammospheres number and size were evaluated using a microscope fitted with a ruler. Random six areas were chosen for quantifying spheres number.

2.4. Cell migration and invasion assay

Cell spheres formed by CC cells were digested and re-suspended, and 8×10^4 cells were added to the upper chamber of 24-well MILLICell Hanging Cell Culture inserts 8 mm PET (MILLIPORE) pre-coated with BD BioCoat Matrigel followed by Tanshinone IIA treatment or not, 800 µl medium containing 20% FBS was used as a chemo-attractant in the bottom chamber. After 24 h for migration and 36 h for invasion, cells migrating and invading into the underside were fixed in methanol for 15 min, and stained with 0.1% viola crystalline solution for 15 min. Five random fields from each of the triplicate were counted by using phase contrast microscopy. Quantification was carried out by measuring with Microplate Reader (OD 570 nm) after being destained with 30% glacial acetic acid.

2.5. Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted using TRNzol (Cat # DP405-02, TIANGEN, Beijing, China) according to the manufacturer's protocols. Then total RNA was reverse transcribed into cDNA using TIANScript RT Kit (Cat # KR104-01, TIANGEN) following the standard protocols. Afterwards, mRNA expression was examined using Quant qRT-PCR kit (SYBR Green) (Cat # FP302-01, TIANGEN) and carried out on an ABI Prism 7500 Detection System (Applied Biosystems, Inc., USA). GAPDH was used as an internal reference. mRNA expression was measured using $2^{-\Delta\Delta Ct}$ method.

2.6. Western blot

The detailed procedure was referred to the previous study [16]. The primary antibodies against HuR (ab200342), YAP (ab52771), ALDH1 (ab9883), Cleaved caspase 3 (ab2302), Cleaved PARP1 (ab32064), Nanog (ab80892) and CTGF (ab6992) were purchased from Abcam. The primary antibodies against Caspase 3 (Cat # 19677-1-AP), PARP1 (Cat # 13371-AP), E-cadherin (Cat 20874-1-AP) and Vimentin (Cat # 10366-1-AP) were purchased from proteintech, GAPDH primary antibody (Cat # AF1186) was purchased from Beyotime (Beijing, China). Secondary peroxidase-conjugated goat anti-rabbit (Cat # A0208) and Peroxidase-conjugated goat anti-mouse (A0216) were purchased from Beyotime. Immunoblots were exposed using the ECL Plus (Solarbio Life Sciences, Beijing, China), following manufacturer's instructions using IVIS-Lumina imaging system (Caliper Life Sciences, MA, USA).

2.7. RNA binding protein immunoprecipitation (RIP) assay

The detailed procedure was mentioned before [17]. Briefly, CC cells were lysed with 25 mM Tris–HCl buffer (pH 7.5) and 100 U/ml RNase inhibitor (Sigma), and then incubated with protein-A Sepharose beads pre-coated with 3 µg anti-HuR (ab200342) antibody or control rabbit IgG for 1.5 h at 4 °C. The RNA-protein complexes were pulled-down by protein A/G agarose beads and RNA was extracted with TRNzol, then YAP mRNA level was detected via qRT-PCR assay.

2.8. Immunofluorescent assay

Cells were cultured on glass bottom plates and treated with Tanshinone IIA. After 24 h, cells were washed with PBS, and then fixed with 4% paraformaldehyde for 15 min, followed by the permeabilization with 0.1% Triton X-100 for 10 min and blocked with 5% BSA in PBS for 1 h at room temperature. Then cells were incubated with antibody against HuR (ab200342) overnight at 4 °C, and washed with PBS for 5 min three times and incubated with FITC-conjugated secondary antibody (Cat # A0562, Beyotime) for 1 h at room temperature, and washed with PBS for three times again and observed under the confocal microscopy.

2.9. Luciferase reporter assay

YAP transcriptional activity was assayed via luciferase reporter assay in cells with different treatment. Briefly, 8xGTIIc-luciferase plasmid was co-transfected into CC cells with β-gal (Ambion, USA) plasmid using Lipofectamine™2000 followed by Tanshinone IIA treatment for 72 h. After that, the luciferase activity of 8xGTIIc-luciferase plasmid was measured using a Luciferase Reporter Assay Kit (cat. no. K801-200, BioVision, Inc., Milpitas, CA, USA). β-gal activity was determined using a β-Galactosidase Enzyme Assay System with Reporter Lysis Buffer (cat. no. E2000, Promega Corporation) following the manufacturer's protocols, and was used as a normalization control for luciferase activity.

2.10. Cell viability assay

Cell spheres were digested and 3×10^3 cells were seeded into 96-well plates, followed by the treatment of adriamycin plus Tanshinone IIA or not, after 24 h, 48 h and 72 h, cell viability was examined by CCK8 assay kit (Cat # HY-K0301, MedChemExpress, USA) according to the manufacturer's recommendation.

2.11. Statistical analysis

All data were expressed as mean ± SD. Means were compared using Student's unpaired test. $P < 0.05$ was considered statistically significant. All experiments were repeated at least three times.

Download English Version:

<https://daneshyari.com/en/article/8525295>

Download Persian Version:

<https://daneshyari.com/article/8525295>

[Daneshyari.com](https://daneshyari.com)