

## Author's Accepted Manuscript

A major component of vitamin E,  $\alpha$ -tocopherol inhibits the anti-tumor activity of crizotinib against cells transformed by EML4-ALK

Yuki Uchihara, Takayuki Kidokoro, Kenji Tago, Tadahiko Mashino, Hiroomi Tamura, Megumi Funakoshi-Tago



[www.elsevier.com/locate/ejphar](http://www.elsevier.com/locate/ejphar)

PII: S0014-2999(18)30093-1  
DOI: <https://doi.org/10.1016/j.ejphar.2018.02.012>  
Reference: EJP71665

To appear in: *European Journal of Pharmacology*

Received date: 22 November 2017  
Revised date: 9 February 2018  
Accepted date: 9 February 2018

Cite this article as: Yuki Uchihara, Takayuki Kidokoro, Kenji Tago, Tadahiko Mashino, Hiroomi Tamura and Megumi Funakoshi-Tago, A major component of vitamin E,  $\alpha$ -tocopherol inhibits the anti-tumor activity of crizotinib against cells transformed by EML4-ALK, *European Journal of Pharmacology*, <https://doi.org/10.1016/j.ejphar.2018.02.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**A major component of vitamin E,  $\alpha$ -tocopherol inhibits the anti-tumor activity of crizotinib against cells transformed by EML4-ALK**

Yuki Uchihara<sup>1</sup>, Takayuki Kidokoro<sup>1</sup>, Kenji Tago<sup>2</sup>, Tadahiko Mashino<sup>3</sup>, Hiroomi Tamura<sup>1</sup>, Megumi Funakoshi-Tago<sup>1,\*</sup>

<sup>1</sup>Division of Hygienic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan

<sup>2</sup>Division of Structural Biochemistry, Department of Biochemistry, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke-shi, Tochigi-ken 329-0498, Japan

<sup>3</sup>Division of Medicinal Chemistry and Bio-organic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan

**\*Corresponding author:** Division of Hygienic Chemistry, Faculty of Pharmacy, Keio University, 1-5-30 Shibakoen, Minato-ku, Tokyo 105-8512, Japan; Tel.: +81-3-5400-2689, fax: +81-3-5400-2689; tago-mg@pha.keio.ac.jp

**Abstract**

Crizotinib is an inhibitor of anaplastic lymphoma kinase (ALK) and is of significant therapeutic benefit to patients with non-small cell lung cancer (NSCLC) harboring the *EML4-ALK* fusion gene. In the present study, we demonstrated that  $\alpha$ -tocopherol, a major component of vitamin E, attenuated the effects of crizotinib independently of its

Download English Version:

<https://daneshyari.com/en/article/8529201>

Download Persian Version:

<https://daneshyari.com/article/8529201>

[Daneshyari.com](https://daneshyari.com)