

Phenolic acid and flavonoid-rich fraction of *Sasa quelpaertensis* Nakai leaves prevent alcohol induced fatty liver through AMPK activation

Kalahe Hewage Iresha Nadeeka Madushani Herath, Jinhee Cho, Areum Kim, Tae Kil Eom, Ju-Sung Kim, Jae-Bum Kim, Yang Hoi Doh, Youngheun Jee



PII: S0378-8741(18)31288-1
DOI: <https://doi.org/10.1016/j.jep.2018.06.008>
Reference: JEP11397

To appear in: *Journal of Ethnopharmacology*

Received date: 11 April 2018

Revised date: 8 June 2018

Accepted date: 11 June 2018

Cite this article as: Kalahe Hewage Iresha Nadeeka Madushani Herath, Jinhee Cho, Areum Kim, Tae Kil Eom, Ju-Sung Kim, Jae-Bum Kim, Yang Hoi Doh and Youngheun Jee, Phenolic acid and flavonoid-rich fraction of *Sasa quelpaertensis* Nakai leaves prevent alcohol induced fatty liver through AMPK activation, *Journal of Ethnopharmacology*, <https://doi.org/10.1016/j.jep.2018.06.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Phenolic acid and flavonoid-rich fraction of *Sasa quelpaertensis* Nakai leaves prevent alcohol induced fatty liver through AMPK activation

Kalahe Hewage Iresha Nadeeka Madushani Herath^a, Jinhee Cho^b, Areum Kim^a, Tae Kil Eom^c, Ju-Sung Kim^d, Jae-Bum Kim^e, Yang Hoi Doh^f, Youngheun Jee^{a,b*}

^aInterdisciplinary Graduate Program in Advanced Convergence Technology & Science, Jeju National University, Jeju 63243, Republic of Korea

^bDepartment of Veterinary Medicine and Veterinary Medical Research Institute, Jeju National University, Jeju 63243, Republic of Korea

^cSubtropical/Tropical Organism Gene Bank, SARI, Jeju National University, Jeju 63243, Korea

^dMajors in Plant Resource and Environment, College of Agriculture and Life Sciences, SARI, Jeju National University, Jeju 63243, Republic of Korea

^eDepartment of Systems Management Engineering, Sungkyunkwan University, Suwon 440-746, Republic of Korea

^fDepartment of Electronic Engineering, Jeju National University, Jeju 63243, Republic of Korea
madushaniherath001@gmail.com

joejh89@nate.com

orange5687@naver.com

taekil7@hanmail.net

aha2011@jejunu.ac.kr

kjbnhg@kitech.re.kr

yhdoh@jejunu.ac.kr

yhjee@jejunu.ac.kr

* Corresponding author. Address: Department of Veterinary Medicine and Veterinary Medical Research Institute, Jeju National University, 102 Jejudaehakno, Jeju 63243, Republic of Korea. Tel.: +82 64 754 3374; Fax: +82 64 756 3354

Abstract

Ethnopharmacological relevance

Sasa quelpaertensis Nakai is an edible dwarf bamboo cultivated mainly in Jeju Island, South Korea and its leaf displays various health-promoting properties including antioxidant scavenging.

Download English Version:

<https://daneshyari.com/en/article/8532158>

Download Persian Version:

<https://daneshyari.com/article/8532158>

[Daneshyari.com](https://daneshyari.com)