


## Accepted Manuscript

The mechanisms behind decreased internalization of angiotensin II type 1 receptor

Jingwei Bian, Suli Zhang, Ming Yi, Mingming Yue, Huirong Liu



PII: S1537-1891(17)30304-X  
DOI: doi:[10.1016/j.vph.2018.01.008](https://doi.org/10.1016/j.vph.2018.01.008)  
Reference: VPH 6451  
To appear in: *Vascular Pharmacology*  
Received date: 12 October 2017  
Revised date: 19 January 2018  
Accepted date: 24 January 2018

Please cite this article as: Jingwei Bian, Suli Zhang, Ming Yi, Mingming Yue, Huirong Liu , The mechanisms behind decreased internalization of angiotensin II type 1 receptor. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Vph*(2018), doi:[10.1016/j.vph.2018.01.008](https://doi.org/10.1016/j.vph.2018.01.008)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The mechanisms behind decreased internalization of angiotensin II type 1 receptor

Jingwei Bian<sup>a</sup>, Suli Zhang<sup>a</sup>, Ming Yi<sup>a</sup>, Mingming Yue<sup>a</sup>, Huirong Liu<sup>a,b,\*</sup>

<sup>a</sup>Department of Physiology & Pathophysiology, School of Basic Medical Sciences, Capital Medical University, Beijing, China;

<sup>b</sup>Beijing Key Laboratory of Cardiovascular Diseases and Related Metabolic Dysfunction, Capital Medical University, Beijing, China;

**\*Address for correspondence:**

Huirong Liu, PhD

Department of Physiology & Pathophysiology, School of Basic Medical Sciences, Capital Medical University 10 Xitoutiao, You An Men Street, Beijing City, 100069, China

Phone: +86-10-83911830

E-mail: liuhr2000@126.com

**Abstract**

The internalization of angiotensin II type 1 receptor (AT<sub>1</sub>R) plays an important role in

Download English Version:

<https://daneshyari.com/en/article/8540488>

Download Persian Version:

<https://daneshyari.com/article/8540488>

[Daneshyari.com](https://daneshyari.com)