

Brazilian Journal
of Pharmacognosy

REVISTA BRASILEIRA DE FARMACOGNOSIA

www.elsevier.com/locate/bjp



Original Article

Impact of light quality on flavonoid production and growth of *Hyptis marrubioides* Epling seedlings cultivated *in vitro*

Rita Cassia N. Pedroso^a, Núbia Angelica A. Branquinho^a, Alessandra C.B.A.M. Hara^a, Alan Carlos Costa^a, Fabiano Guimarães Silva^a, Leticia P. Pimenta^b, Marcio Luiz A. Silva^b, Wilson Roberto Cunha^b, Patricia M. Pauletti^b, Ana Helena Januario^{b,*}

^a Instituto Federal de Educação, Ciência e Tecnologia Goiano, Câmpus Rio Verde, Rio Verde, GO, Brazil

^b Núcleo de Pesquisa em Ciências Exatas e Tecnológicas, Universidade de Franca, Franca, SP, Brazil

ARTICLE INFO

Article history:

Received 2 November 2016

Accepted 22 December 2016

Available online xxx

ABSTRACT

Hyptis marrubioides Epling, Lamiaceae, a species from Brazilian Cerrado, has been used against gastrointestinal infections, skin infections, pain, and cramps. Herein, *H. marrubioides* seedlings were cultured *in vitro* under different wavelengths (white, blue, green, red, and yellow) with $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ irradiance and a 16-h photoperiod. After 20 and 30 days of cultivation, shoot length, leaf number, fresh mass, and dry mass were evaluated. The flavonoid rutin content was determined by the HPLC-DAD method. The shoots were longer in plants cultivated under yellow (16.603 ± 0.790 cm, 1.8-fold), red (15.465 ± 0.461 cm, 1.7-fold), and green (14.677 ± 0.737 cm, 1.6-fold) lights than in control plants exposed to white light (9.203 ± 0.388 cm). The number of leaves increased in plants exposed to red (23.425 ± 1.138 , 1.1-fold) and green (22.725 ± 1.814 , 1.1-fold) lights, compared to control plants (20.133 ± 0.827). Fresh (0.665 ± 0.048 g, 1.2-fold) and dry (0.066 ± 0.005 g, 1.3-fold) mass of seedlings were the highest in seedlings grown under red light, compared to seedlings grown under white light (0.553 ± 0.048 and 0.028 ± 0.004 , respectively). However, rutin production was higher under white (0.308 mg g^{-1} of dry weight) and blue lights (0.298 mg g^{-1} of dry weight). Thus, red light induces plant growth and increases leaf number and dry weight in *in vitro*-cultivated *H. marrubioides*, whereas blue and white lights promote the greatest rutin accumulation.

© 2017 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The genus *Hyptis*, Lamiaceae, comprises about 300 species, widely distributed, occurring mainly in tropical regions of the Americas and Africa. *Hyptis marrubioides* Epling, commonly known as mint-of-field, is a species of the Brazilian Cerrado traditionally used to treat gastrointestinal and skin infections, pain, and cramps (McNeil et al., 2011). The pharmacological potential of *H. marrubioides* has been previously investigated, and most studies have focused on examination of the chemical composition of essential oils, the main components of which are the sesquiterpenes caryophylla-4(14),8(15)-dien-5 β -ol, eudesma-4(15),7-dien-1 β -ol, caryophyllene oxide, and β -caryophyllene (Sales et al., 2007; McNeil et al., 2011). Essential oils of *H. marrubioides* have also been studied to investigate their potential for pest control in agriculture,

such as the prevention and control of Asian soybean rust (Silva et al., 2012b) and treatment of *Colletotrichum truncatum*-infected soybean seeds (Silva et al., 2012a).

The use of micropropagation techniques has great advantages in the production of high-quality seedlings, as they improve their pharmacological potential (Rao and Ravishankar, 2002) by ensuring the reproduction of identical, high-quality plants (Serafini et al., 2001), which in turn enhances the biosynthesis of special metabolites (Bhuiyan and Adachi, 2003; Zhao et al., 2005). However, the accumulation of secondary metabolites in plants is influenced by various environmental factors such as light quality, UV irradiation, temperature, irrigation, nutrient deficiency, pathogen attack, and heavy metal stress (Dixon and Paiva, 1995; Winkel-Shirley, 2002; Kopsell and Kopsell, 2008).

Light is a key abiotic elicitor in plants that affects, directly or indirectly, growth and development of plants, mainly of those distributed in high-latitude areas (Morini and Muleo, 2003; OuYang et al., 2015). Plant responses do not depend only on the absence or presence of light but also on the variation in light quality (Felippe,

* Corresponding author.

E-mail: ana.januario@unifran.edu.br (A.H. Januario).

<http://dx.doi.org/10.1016/j.bjp.2016.12.004>

0102-695X/© 2017 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1986). The action of light on plants occurs mainly in two aspects: first, the light source provides the energy required by the plant for photosynthesis, especially red and blue light, and second, as a signal received by photoreceptors it regulates growth, differentiation, and plant metabolism (Wang et al., 2001).

Light is one of the most significant environmental factors affecting the accumulation of flavonoids in plants (Huché-Thélier et al., 2016). However, the biosynthesis of flavonoids in response to light quality need to be better understood, especially in *in vitro* culture.

Studies involving the influence of light on the production of secondary metabolites and growth parameters in *H. marrubioides* seedlings cultivated *in vitro* are non-existent. In a previous study, we have evaluated the phytochemical profile of *H. marrubioides* microplants inoculated with isolates of bacteria and endophytic fungi (Vitorino et al., 2013). Photosynthetic studies in *H. marrubioides in vitro* culture involving measuring gas exchange were also conducted by our research group and the irradiance effects, flow rate, and air humidity parameters were investigated. In addition, the chlorophyll a fluorescence and chloroplastidic pigment content were also assessed (Costa et al., 2014). Hence, the aim of the present study was to assess the effect of light quality on growth of *H. marrubioides* seedlings *in vitro* and their flavonoid production. This study furthers our understanding of the factors that promote the growth and increase the resistance of this medicinal species, through the production of flavonoids.

Materials and methods

Plant material and *in vitro* cultivation

Seedlings previously established *via in vitro* germination of seeds suitable for inoculation obtained on the experimental field of the Laboratory of Plant Tissue Culture of the Instituto Federal de Educação, Ciência e Tecnologia Goiano, Campus Rio Verde. A voucher specimen (HRV71) has been deposited at the Herbarium of the Institute (Herbarium HRV).

The seeds were disinfected with 0.2% Bendazol (carbendazim) and 0.2% Alterno (tebuconazole) for 1 h, followed by treatment with 1% sodium hypochlorite for 30 min, and then rinsed thrice with sterile distilled water. Seeds were germinated and maintained on Murashige & Skoog medium (Murashige and Skoog, 1962) supplemented with 30 g l⁻¹ of sucrose and solidified with agar at 3.5 g l⁻¹; the pH was adjusted to 5.8 before autoclaving. The cultures were incubated in a growth chamber for 30 days at 50 μmol m⁻² s⁻¹ photosynthetically active radiation (PAR), at an average temperature of 23 ± 1 °C, and a photoperiod of 16 h. Thereafter, the seedlings were sub-cultured on the same medium and incubated in a growth cabinet for 10 days at 50 μmol m⁻² s⁻¹ PAR, 23 ± 1 °C, and 16-h photoperiod. All experiments were carried out in glass flasks containing 50 ml of semisolid medium with five seedlings per flask with a total of forty seedlings.

Light conditions

After 10 days, the cultures were transferred to a cabinet and exposed to continuing irradiation of different light spectra: white (300–750 nm), blue (400–490 nm), green (490–560 nm), red (600–700 nm), and yellow (560–590 nm). The specific light conditions were maintained using TP 40W lamps (Taschibra® Indaial, Santa Catarina, Brazil), at a light intensity of 50 μmol m⁻² s⁻¹ and a photoperiod of 16 h. The cultures were evaluated after 20 and 30 days of light exposure. Cultures grown under white light were used as a control. For weight measurements, the whole seedlings without roots was considered.

Chemical analysis

The seedlings were dried at 35 °C in a forced air circulation oven and the dry biomass (200 mg) from the *in vitro* culture of *H. marrubioides* was extracted with 4 ml methanol of high performance liquid chromatography (HPLC) grade using ultrasound bath (30 min). The samples had been previously filtered through a 0.2-μm syringe filter with hydrophilic PTFE membrane (Advantec, Dublin, CA, USA) and transferred into 1-ml HPLC vials. These procedures were conducted in triplicate.

For the quantitative analysis, solutions were done in HPLC grade methanol to obtain solutions containing 0.500, 0.250, 0.125, and 0.063 mg ml⁻¹ rutin. Each standard solution was injected in triplicate. The calibration curve was constructed to determine the linearity of the method by plotting the peak area versus the concentration of the substance in mg ml⁻¹. The quantitative analysis by HPLC with diode-array detection (DAD) was carried out on a Shimadzu Prominence LC-20AD binary system equipped with a DGU-20A5 degasser, an SPD-20A series diode array detector, a CBM-20A communication bus module, an SIL-20A HT autosampler, and a CTO-20A column oven (Shimadzu, Kyoto, Japan). The analyses were conducted on a Gemini ODS column (250 × 4.6 mm, 5 μm; Phenomenex, Aschaffenburg, Germany) equipped with a pre-column with the same material under the following conditions: injection volume set at 20 μl, flow rate of 1.0 ml min⁻¹, mobile phase was CH₃OH/H₂O/HOAc (5:94.9:0.1, v/v/v) delivered in a linear gradient until 100% CH₃OH in 30 min, 10 min at 100% CH₃OH, and 20 min to return to the initial conditions. The UV detection was set at 254 nm and 40 °C.

Methanol used in the experiments was HPLC grade and it was obtained from J. T. Baker (Avantor Performance Materials, Center Valley, PA, USA). Ultrapure water was obtained by passing redistilled water through a Direct-Q UV3 system from Millipore (Billerica, MA, USA). The flavonoid rutin used as external standard was acquired from the standard bank of the Natural Products Group of the Universidade de Franca.

Statistical analyses

The statistical analysis was performed in Sisvar 5.3 software (Ferreira, 2011). The experiment was conducted in a completely randomized design with four replications. The multi-factorial analysis of variance followed by Tukey multiple comparison tests were used for statistical comparisons ($p \leq 0.05$).

Results

Seedlings growth

The measured growth parameters of *H. marrubioides* seedlings cultivated *in vitro* under different wavelengths showed significant difference (Table 1). During the 20-day cultivation, the seedlings exposed to red light showed the highest shoot lengths (10.175 ± 0.669 cm), which was 1.69-fold higher than the corresponding lengths of control seedlings (6.035 ± 0.511 cm). In addition, after 30 days of irradiation with green, red, and yellow lights, the length of the shoots increased 1.8-fold compared to seedlings exposed to white light. The number of expanded leaves was not affected by light conditions after 20 days of treatment; however, after 30 days of culture, green and red light promoted the most the increase in the number of leaves (22.725 ± 1.814 and 23.425 ± 1.138, respectively).

Shoot dry and fresh weights did not differ among the seedlings irradiated with different lights for 20 days. In contrast, after 30 days of illumination with red light, dry biomass of *H. marrubioides*

Download English Version:

<https://daneshyari.com/en/article/8543300>

Download Persian Version:

<https://daneshyari.com/article/8543300>

[Daneshyari.com](https://daneshyari.com)