Accepted Manuscript



Assessment of the impact of aerosol from a potential modified risk tobacco product compared with cigarette smoke on human organotypic oral epithelial cultures under different exposure regimens

Filippo Zanetti, Alain Sewer, Elena Scotti, Bjoern Titz, Walter K. Schlage, Patrice Leroy, Athanasios Kondylis, Gregory Vuillaume, Anita R. Iskandar, Emmanuel Guedj, Keyur Trivedi, Thomas Schneider, Ashraf Elamin, Florian Martin, Nikolai V. Ivanov, Stefan Frentzel, Manuel C. Peitsch, Julia Hoeng

PII: S0278-6915(18)30136-4

DOI: 10.1016/j.fct.2018.02.062

Reference: FCT 9633

To appear in: *Food and Chemical Toxicology*

Received Date: 29 December 2017

Revised Date: 20 February 2018

Accepted Date: 27 February 2018

Please cite this article as: Zanetti, F., Sewer, A., Scotti, E., Titz, B., Schlage, W.K., Leroy, P., Kondylis, A., Vuillaume, G., Iskandar, A.R., Guedj, E., Trivedi, K., Schneider, T., Elamin, A., Martin, F., Ivanov, N.V., Frentzel, S., Peitsch, M.C., Hoeng, J., Assessment of the impact of aerosol from a potential modified risk tobacco product compared with cigarette smoke on human organotypic oral epithelial cultures under different exposure regimens, *Food and Chemical Toxicology* (2018), doi: 10.1016/j.fct.2018.02.062.

CHTP1.2 aerosol OR 3R4F Smoke

Buccal Organotypic cultures





Gingival Organotypic cultures





Sample collection

CHTP1.2 Relative Effect Reduction compared to 3R4F (%)