Accepted Manuscript



T-2 toxin-induced toxicity in neuroblastoma-2a cells involves the generation of reactive oxygen, mitochondrial dysfunction and inhibition of Nrf2/HO-1 pathway

Xiya Zhang, Ying Wang, Tony Velkov, Shusheng Tang, Chongshan Dai

PII: S0278-6915(18)30075-9

DOI: 10.1016/j.fct.2018.02.010

Reference: FCT 9581

To appear in: *Food and Chemical Toxicology*

Received Date: 28 August 2017

Revised Date: 17 January 2018

Accepted Date: 6 February 2018

Please cite this article as: Zhang, X., Wang, Y., Velkov, T., Tang, S., Dai, C., T-2 toxin-induced toxicity in neuroblastoma-2a cells involves the generation of reactive oxygen, mitochondrial dysfunction and inhibition of Nrf2/HO-1 pathway, *Food and Chemical Toxicology* (2018), doi: 10.1016/j.fct.2018.02.010.

T-2 toxin-induced toxicity in neuroblastoma-2a cells involves the generation of reactive oxygen, mitochondrial dysfunction and inhibition of Nrf2/HO-1 pathway

Xiya Zhang[1,2], Ying Wang[2], Tony Velkov[3], Shusheng Tang[2*], Chongshan Dai[2*]

[1]College of Food Science and Technology, Henan Agricultural University, 63 Nongye Road, Zhengzhou 450002, P. R. China. [2]College of Veterinary Medicine, China Agricultural University, Beijing 100193, P. R. China. [3]Department of Pharmacology & Therapeutics, School of Biomedical Sciences, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Parkville, VIC, 3010, Australia.

Running title: T-2 toxin induces apoptosis inhibits Nrf2/HO-1 pathway

*Joint corresponding authors:

Chongshan Dai. Telephone: +86 10 6273 3377; Fax: +86 10 6273 1032. Telephone:

E-mail: daichongshan@163.com

Shusheng Tang. Telephone: +86 10 6273 3857; Fax: +86 10 6273 1032. E-mail: tssfj@163.com