


Accepted Manuscript

Title: TLR2 and TLR3 expression as a biomarker for the risk of doxorubicin-induced heart failure

Authors: Shao Liang, Cai Xinyong, Zhu Hongmin, Wang Jing, Hong Lang, Zhang Ping



PII: S0378-4274(18)31469-3
DOI: <https://doi.org/10.1016/j.toxlet.2018.06.1219>
Reference: TOXLET 10255

To appear in: *Toxicology Letters*

Received date: 14-5-2018
Revised date: 19-6-2018
Accepted date: 26-6-2018

Please cite this article as: Liang S, Xinyong C, Hongmin Z, Jing W, Lang H, Ping Z, TLR2 and TLR3 expression as a biomarker for the risk of doxorubicin-induced heart failure, *Toxicology Letters* (2018), <https://doi.org/10.1016/j.toxlet.2018.06.1219>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

TLR2 and TLR3 expression as a biomarker for the risk of doxorubicin-induced heart failure

Shao Liang^{1, 2*}, Cai Xinyong¹, Zhu Hongmin¹, Wang Jing¹, Hong Lang^{1, 2*}, Zhang Ping^{3*}

¹ Department of Cardiology, Jiangxi Provincial People's Hospital, No. 92 Aiguo Road, Donghu District, Nanchang, 330006, Jiangxi, People's Republic of China.

² Jiang Xi provincial Institute of Cardiovascular Diseases, No. 92 Aiguo Road, Donghu District, Nanchang, 330006, Jiangxi, People's Republic of China.

³ Department of Neurology, Jiangxi Provincial People's Hospital, No. 92 Aiguo Road, Donghu District, Nanchang, 330006, Jiangxi, People's Republic of China.

* **Corresponding author:** Shao Liang, E-mail: shaoliang021224@hotmail.com; Zhang Ping, E-mail: zhangkiki520@126.com; Hong Lang, E-mail: honglang13@163.com

Highlights

- doxorubicin is believed to cause cardiotoxicity by inflammation.
- Toll like receptors play important roles in doxorubicin-induced cardiotoxicity
- TLR2 and TLR3 expression as a biomarker for the risk of doxorubicin-induced heart failure

Download English Version:

<https://daneshyari.com/en/article/8553101>

Download Persian Version:

<https://daneshyari.com/article/8553101>

[Daneshyari.com](https://daneshyari.com)