Accepted Manuscript



ROS-mediated oligomerization of VDAC2 is associated with quinocetone-induced apoptotic cell death

Xiayun Yang, Shusheng Tang, Daowen Li, Bin Li, Xilong Xiao

PII: S0887-2333(17)30371-5
DOI: doi:10.1016/j.tiv.2017.12.005
Reference: TIV 4182

To appear in: *Toxicology in Vitro*

Received date: 15 August 2017
Revised date: 13 October 2017
Accepted date: 5 December 2017

Please cite this article as: Xiayun Yang, Shusheng Tang, Daowen Li, Bin Li, Xilong Xiao , ROS-mediated oligomerization of VDAC2 is associated with quinocetone-induced apoptotic cell death. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Tiv(2017), doi:10.1016/j.tiv.2017.12.005

ROS-mediated oligomerization of VDAC2 is associated with quinocetone-induced apoptotic cell death

Xiayun Yang[a], Shusheng Tang[a], Daowen Li[a], Bin Li[a], Xilong Xiao[a*]

[a] Department of Pharmacology and Toxicology, College of Veterinary Medicine, China Agricultural University, Yuanminyuan West Road No. 2, Beijing, Haidian District 100193, China

*Corresponding author: Xilong Xiao. Tel: +86 1062733377; Fax: +86 10 6273 1032; E-mail address: xiaoxl@cau.edu.cn