


Accepted Manuscript

Rebamipide suppresses 5-fluorouracil-induced cell death via the activation of Akt/mTOR pathway and regulates the expression of Bcl-2 family proteins

Masanobu Tsubaki, Tomoya Takeda, Ryo-ta Asano, Tomoyuki Matsuda, Shin-ichiro Fujimoto, Tatsuki Itoh, Motohiro Imano, Takao Satou, Shozo Nishida



PII: S0887-2333(17)30316-8
DOI: doi:[10.1016/j.tiv.2017.10.019](https://doi.org/10.1016/j.tiv.2017.10.019)
Reference: TIV 4150
To appear in: *Toxicology in Vitro*
Received date: 15 April 2017
Revised date: 12 September 2017
Accepted date: 16 October 2017

Please cite this article as: Masanobu Tsubaki, Tomoya Takeda, Ryo-ta Asano, Tomoyuki Matsuda, Shin-ichiro Fujimoto, Tatsuki Itoh, Motohiro Imano, Takao Satou, Shozo Nishida , Rebamipide suppresses 5-fluorouracil-induced cell death via the activation of Akt/mTOR pathway and regulates the expression of Bcl-2 family proteins. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Tiv*(2017), doi:[10.1016/j.tiv.2017.10.019](https://doi.org/10.1016/j.tiv.2017.10.019)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Rebamipide suppresses 5-fluorouracil-induced cell death via the activation of Akt/mTOR pathway and regulates the expression of Bcl-2 family proteins

Running title: Rebamipide suppresses 5-FU-induced cell death

Masanobu Tsubaki¹, Tomoya Takeda¹, Ryo-ta Asano¹, Tomoyuki Matsuda¹, Shin-ichiro Fujimoto¹, Tatsuki Itoh², Motohiro Imano³, Takao Satou⁴, and Shozo Nishida^{1,*}

¹Division of Pharmacotherapy, Kindai University Faculty of Pharmacy, Kowakae, Higashi-Osaka, Japan; ²Department of Food Science and Nutrition, Kindai University Faculty of Agriculture, Nara, Nara, Japan.; ³Department of Surgery, Kindai University Faculty of Medicine, Osakasayama, Osaka, Japan.; ⁴Department of Pathology, Kindai University Faculty of Medicine, Osakasayama, Osaka, Japan.

Keywords: rebamipide; oral mucositis; 5-FU; Akt; mTOR

*Corresponding author: Shozo Nishida, Ph.D. Postal address: Division of Pharmacotherapy, Kindai University Faculty of Pharmacy, Kowakae, Higashi-Osaka 577-8502, Japan. Tel.: +81-6-6721-2332 ex.3851; Fax +81-6-6730-1394; E-mail address: nishida@phar.kindai.ac.jp

Download English Version:

<https://daneshyari.com/en/article/8554126>

Download Persian Version:

<https://daneshyari.com/article/8554126>

[Daneshyari.com](https://daneshyari.com)