


Accepted Manuscript

Legume consumption is inversely associated with type 2 diabetes incidence in adults: a prospective assessment from the PREDIMED study

Nerea Becerra-Tomás, RD, Andrés Díaz-López, PhD, Núria Rosique-Esteban, RD, Emilio Ros, PhD, Pilar Buil-Cosiales, PhD, Dolores Corella, PhD, Ramon Estruch, PhD, Montserrat Fitó, PhD, Lluís Serra-Majem, PhD, Fernando Arós, PhD, Rosa Maria Lamuela-Raventós, PhD, Miquel Fiol, PhD, José Manuel Santos-Lozano, PhD, Javier Diez-Espino, PhD, Olga Portoles, PhD, Jordi Salas-Salvadó, PhD



PII: S0261-5614(17)30106-1

DOI: [10.1016/j.clnu.2017.03.015](https://doi.org/10.1016/j.clnu.2017.03.015)

Reference: YCLNU 3086

To appear in: *Clinical Nutrition*

Received Date: 8 November 2016

Revised Date: 17 February 2017

Accepted Date: 14 March 2017

Please cite this article as: Becerra-Tomás N, Díaz-López A, Rosique-Esteban N, Ros E, Buil-Cosiales P, Corella D, Estruch R, Fitó M, Serra-Majem L, Arós F, Lamuela-Raventós RM, Fiol M, Santos-Lozano JM, Diez-Espino J, Portoles O, Salas-Salvadó J, PREDIMED Study Investigators, Legume consumption is inversely associated with type 2 diabetes incidence in adults: a prospective assessment from the PREDIMED study, *Clinical Nutrition* (2017), doi: 10.1016/j.clnu.2017.03.015.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Legume consumption is inversely associated with type 2 diabetes incidence in adults: a prospective assessment from the PREDIMED study

Nerea Becerra-Tomás RD^{a,b}, Andrés Díaz-López PhD^{a,b}, Núria Rosique-Esteban RD^a, Emilio Ros PhD^{b,c}, Pilar Buil-Cosiales PhD^{b,d}, Dolores Corella PhD^{b,e}, Ramon Estruch PhD^{b,f}, Montserrat Fitó PhD^{b,g}, Lluís Serra-Majem PhD^{b,h}, Fernando Arós PhD^{b,i}, Rosa Maria Lamuela-Raventós PhD^{b,j}, Miquel Fiol PhD^{b,k}, José Manuel Santos-Lozano PhD^{b,l}, Javier Diez-Espino PhD^{b,d}, Olga Portoles PhD^{b,e} and Jordi Salas-Salvadó PhD^{a,b*}; PREDIMED Study Investigators*

^aHuman Nutrition Unit, University Hospital of Sant Joan de Reus, Department of Biochemistry and Biotechnology, Faculty of Medicine and Health Sciences, Institut d'Investigació Sanitària Pere Virgili (IISPV), Rovira i Virgili University, Reus, Spain.

^bCentro de Investigación Biomédica en Red Fisiopatología de la Obesidad y la Nutrición (CIBEROBN), Institute of Health Carlos III, Madrid, Spain. ^cLipid Clinic, Endocrinology and Nutrition Service, IDIBAPS, Hospital Clinic, University of Barcelona, Barcelona, Spain. ^dAtención Primaria, Servicio Navarro de Salud- Osasunbidea, 31010 Navarra, Spain and IdiSNA, Navarra Institute for Health Research.

^eDepartment of Preventive Medicine, University of Valencia, Valencia, Spain.

^fDepartment of Internal Medicine, August Pi i Sunyer Institute of Biomedical Research (IDIBAPS), Hospital Clinic, University of Barcelona, Barcelona, Spain.

^gCardiovascular Risk and Nutrition Research Group, Institut Hospital del Mar d'Investigacions Mèdiques, Barcelona Biomedical Research Park, Barcelona, Spain.

^hDepartment of Clinical Sciences, University of Las Palmas de Gran Canaria, Las Palmas, Spain. ⁱDepartment of Cardiology, University Hospital Araba, Vitoria, Spain.

Download English Version:

<https://daneshyari.com/en/article/8586650>

Download Persian Version:

<https://daneshyari.com/article/8586650>

[Daneshyari.com](https://daneshyari.com)