

Accepted Manuscript

Increase in tear film lipid layer thickness after instillation of 3% diquafosol ophthalmic solution in healthy human eyes

Shima Fukuoka, MD, Reiko Arita, MD, PhD

PII: S1542-0124(16)30288-9

DOI: [10.1016/j.jtos.2017.03.005](https://doi.org/10.1016/j.jtos.2017.03.005)

Reference: JTOS 226

To appear in: *Ocular Surface*

Received Date: 29 December 2016

Revised Date: 20 March 2017

Accepted Date: 20 March 2017

Please cite this article as: Fukuoka S, Arita R, Increase in tear film lipid layer thickness after instillation of 3% diquafosol ophthalmic solution in healthy human eyes, *Ocular Surface* (2017), doi: 10.1016/j.jtos.2017.03.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

SECTION: Original Research, Ali Djalilian, MD, Editor

TITLE: Increase in Tear Film Lipid Layer Thickness after Instillation of 3% Diquafosol Ophthalmic Solution in Healthy Human Eyes

AUTHORS: Shima Fukuoka, MD,¹⁻³ and Reiko Arita, MD, PhD²⁻⁴

SHORT TITLE: Diquafosol and tear film lipid layer thickness/ FUKUOKA AND ARITA

FOOTNOTES

Accepted for publication March 2017.

From ¹Omiya Hamada Eye Clinic, Saitama, ²Lid and Meibomian Gland Working Group (LIME),

³Department of Ophthalmology, University of Tokyo School of Medicine, Tokyo, Japan, and

⁴Department of Ophthalmology, Itoh Clinic, Saitama, Japan.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Dr. Arita has received personal fees from Kao Corporation (Tokyo, Japan) and Japan Focus Company Ltd. (Tokyo, Japan), has received financial support from TearScience (Morrisville, NC, USA), and holds patents on the noncontact meibography system used in the study (JP patent registration no. 5281846, U.S. patent publication no. 2011-0273550A1, EP patent publication no. 2189108A1). Dr. Fukuoka declares no potential conflict of interest.

Corresponding author: Reiko Arita, MD, PhD, Department of Ophthalmology, Itoh Clinic, 626-11 Minami-Nakano, Minuma-ku, Saitama, Saitama 337-0042, Japan. Tel.: +81-48-686-5588 Fax: +81-48-686-8485 Email: ritoh@za2.so-net.ne.jp

Download English Version:

<https://daneshyari.com/en/article/8591239>

Download Persian Version:

<https://daneshyari.com/article/8591239>

[Daneshyari.com](https://daneshyari.com)