ARTICLE IN PRESS

REV BRAS ORTOP, 2018:xxx(xx):xxx-xxx







Case Report

Tibial tubercle fracture associated with distal rupture of the patellar tendon: case report[☆]

André Lourenço Pereira*, Ângelo Ribeiro Vaz de Faria, Túlio Vinícius de Oliveira Campos, Marco Antônio Percope de Andrade, Guilherme Moreira de Abreu e Silva

Hospital das Clínicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

ARTICLE INFO

Article history: Received 12 February 2017 Accepted 11 April 2017 Available online xxx

Keywords: Avulsion fractures Tibial tubercle Rupture Patellar tendon

Palavras-chave: Fraturas avulsão Fraturas da tíbia Ruptura Tendão patelar

ABSTRACT

Avulsion of the tibial tubercle is an uncommon injury that occurs in the young athlete, resulting from an eccentric contraction of the extensor mechanism with the leg fixed to the ground. Concomitant injuries to the patellar tendon are very rare, with few cases reported in the literature. The authors present a case of a 15-year-old basketball player who suffered an avulsion of the tibial tubercle associated with complete distal rupture of the patellar tendon while training. It was treated with open reduction of the tibial fragment and reconstruction of the patellar tendon with mini-anchors and cannulated screws, as well as hamstring autograft tendon reinforcement. The patient showed excellent results and returned to sports after 12 months of follow-up.

© 2018 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Ortopedia e Traumatologia. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Fratura da tuberosidade anterior da tíbia associada à ruptura distal do tendão patelar: relato de caso

RESUMO

A fratura-avulsão da tuberosidade anterior da tíbia é uma lesão incomum que ocorre no jovem atleta, resultado de uma contração excêntrica do mecanismo extensor do joelho com o membro inferior fixo ao solo. Lesões concomitantes ao tendão patelar são muito raras, com poucos casos relatados na literatura. Os autores apresentam o caso de um atleta de basquete de 15 anos que sofreu uma fratura-avulsão da tuberosidade anterior da tíbia associada à ruptura completa distal do tendão patelar durante movimento de arremesso no treino esportivo. O paciente foi tratado com redução aberta da fratura e reparo do tendão patelar com miniâncoras e parafuso poste com reforço tendinoso

E-mail: andrelpereira@live.com (A.L. Pereira).

https://doi.org/10.1016/j.rboe.2018.05.009

2255-4971/© 2018 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Ortopedia e Traumatologia. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Please cite this article in press as: Pereira AL, et al. Tibial tubercle fracture associated with distal rupture of the patellar tendon: case report. Rev Bras Ortop. 2018. https://doi.org/10.1016/j.rboe.2018.05.009

[🌣] Study conducted at Universidade Federal de Minas Gerais, Hospital das Clínicas, Belo Horizonte, MG, Brazil.

^{*} Corresponding author.

REV BRAS ORTOP. 2018; xxx(xx): xxx-xxx

com enxerto autólogo de semitendíneo. O paciente apresentou ótimos resultados e retornou ao esporte após 12 meses de acompanhamento.

© 2018 Publicado por Elsevier Editora Ltda. em nome da Sociedade Brasileira de Ortopedia e Traumatologia. Este é um artigo Open Access sob a licença de CC BY-NC-ND (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Introduction

Avulsion fracture of the anterior tibial tuberosity is a rare injury in young athletes, accounting for less than 1% of all physeal injuries. The fracture mechanism is an eccentric contraction of the quadriceps at the beginning of a jump or during landing. The association with complete distal rupture of the patellar tendon is even rarer, with few cases reported in the literature and no known estimated incidence.

This injury, first classified by Ogden et al., was modified by Frankl et al.² in 1990 to include associated rupture of the patellar tendon. The management of this injury is typically surgical, and a high incidence of preoperative suspicion is necessary to identify the associated rupture of the patellar tendon. The authors present a rare case of anterior tibial tuberosity fracture with distal avulsion of the patellar tendon that occurred in a 15-year-old basketball player, who was surgically treated and presented good postoperative evolution.

Case report

A 15-year-old basketball player presented a left knee injury after a jump while pitching during basketball practice. He complained of severe pain and the impossibility of weight-bearing immediately after the trauma. During the initial evaluation, he presented massive hemarthrosis, a flexion attitude, and inability to actively extend the knee. The initial radiographs indicated a fracture-avulsion of the anterior tibial tuberosity and patella alta (Fig. 1). Magnetic resonance imaging showed a complete distal rupture of the patellar tendon associated with the fracture (Fig. 2).

The patient was admitted to hospital and underwent open reduction and internal fixation of the fracture, and repair of the tendon injury. An anterior longitudinal incision was chosen; the anatomical reduction was secured with two cannulated screws (4.5 mm, Synthes[®]; Fig. 3A). The patellar tendon then underwent repair + grafting with a semitendinosus graft and fixation with two mini-anchors (5 mm, Metabio[®]) and a distal pole screw (4.5 mm, Synthes[®]; Fig. 3B).

Due to the patient's physical size and his high functional demand, the graft was reinforced with autologous semitendinous graft (Kelikian et al.³ technique). After surgery, the knee was immobilized with a brace in extension; the patient was discharged the following day. He was encouraged to perform passive assisted movements until 60 degrees of flexion and kept on crutches for partial weight support. The brace was removed after six weeks; physical therapy was then initiated for strength gain. Total weight-bearing was authorized as tolerated. After 14 weeks, full mobility and radiographic consolidation were observed. Strengthening exercises, through



Fig. 1 – Lateral view radiographs of the knee indicating anterior tibial tuberosity fracture and patella alta.



Fig. 2 – Magnetic resonance imaging demonstrating total patellar tendon rupture.

Download English Version:

https://daneshyari.com/en/article/8599783

Download Persian Version:

https://daneshyari.com/article/8599783

<u>Daneshyari.com</u>