



Original Article

Efficacy evaluation of a protocol for safe hip surgery (total hip arthroplasty)[☆]



Antônio Augusto Guimarães Barros*, Carlos Henrique Cardoso Mendes, Eduardo Frois Temponi, Lincoln Paiva Costa, Carlos Cesar Vassalo, Euler de Carvalho Guedes

Hospital Madre Teresa, Belo Horizonte, MG, Brazil

ARTICLE INFO

Article history:

Received 10 November 2016

Accepted 26 January 2017

Available online 18 August 2017

Keywords:

Hip arthroplasty

Protocols

Postoperative complications

Hospitalization

ABSTRACT

Objective: To propose a multidisciplinary protocol to standardize the care of patients undergoing total hip arthroplasty (THA) and evaluate its effectiveness after implementation.

Methods: Retrospective evaluation of 95 consecutive patients undergoing THA divided into two groups, one group of 47 patients operated before the protocol implementation and 48 after.

Results: Assessing the re-admission rate, among 47 patients evaluated prior to implementation of the protocol, seven (14.9%) were re-admitted, and when observing the 48 patients evaluated after implementation, one (2.1%) was re-admitted, showing statistical significance ($p < 0.05$). The chance of re-admission before the protocol was eight times the chance of hospitalization after implementation (95% CI: 1.01 to 377.7). By comparing the clinical complications among the groups, it was observed that there was a lower rate of complications following implementation of the protocol ($p = 0.006$).

Conclusion: The introduction of a multidisciplinary protocol to standardize the management of patients undergoing THA decreased the rates of rehospitalization and clinical complications after the procedure.

© 2017 Sociedade Brasileira de Ortopedia e Traumatologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Avaliação da eficácia do protocolo para cirurgia segura do quadril (artroplastia total)

RESUMO

Objetivo: Propor um protocolo multidisciplinar para padronização do cuidado dos pacientes que serão submetidos a artroplastia total do quadril (ATQ) e avaliar sua eficácia após a implantação.

Métodos: Avaliação retrospectiva dos resultados de 95 pacientes consecutivos submetidos a ATQ divididos em dois grupos, um com 47 operados antes da implantação do protocolo e 48 após.

Palavras-chave:

Artroplastia de quadril

Protocolos

Complicações pós-operatórias

Hospitalização

[☆] Study conducted at Hospital Madre Teresa, Belo Horizonte, MG, Brazil.

* Corresponding author.

E-mail: antonioagbarros@gmail.com (A.A. Barros).

<http://dx.doi.org/10.1016/j.rboe.2017.08.004>

2255-4971/© 2017 Sociedade Brasileira de Ortopedia e Traumatologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Resultados: Na avaliação da taxa de reinternação, tem-se que entre os 47 pacientes avaliados antes da implantação do protocolo, sete (14,9%) foram reinternados e dos 48 avaliados depois da implantação, um (2,1%) foi reinternado, mostrou-se significância estatística ($p < 0,05$). A chance de reinternação antes da implantação foi oito vezes maior do que a chance de internação após a implantação (IC 95%: 1,01 a 377,7). Ao comparar as complicações clínicas entre os grupos observou-se que houve menor taxa de complicações após a implantação do protocolo ($p = 0,006$).

Conclusão: A introdução de um protocolo multidisciplinar para padronização do manejo do paciente submetido a ATQ diminuiu as taxas de reinternação e de complicações clínicas após o procedimento.

© 2017 Sociedade Brasileira de Ortopedia e Traumatologia. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

For roughly 50 years, total hip arthroplasty (THA) has been one of the most effective orthopedic interventions from a functional and economic standpoint.¹⁻⁴ It is an elective procedure when treating hip arthrosis, and it is performed after careful preoperative evaluation to minimize risks. Advances in tribology, surgical, anesthetic, and rehabilitation techniques have improved the end result. This evolution has led to an increase in surgical indications and in the safety of the procedure,¹ with an overall complication rate of less than 4% and overall 90-day mortality of less than 1%.⁵⁻⁷

Despite these advances, patients undergoing this type of procedure are subject to complications such as infection, implant dislocation, deep venous thrombosis (DVT), and pulmonary embolism (PE), among others,⁵ resulting in increased hospital length of stay and rates of rehospitalization.⁸

Identifying the risk factors for complications as well as applying scientifically effective methods for their prevention are important steps in the strategy to reduce such events, reducing risks to the surgeon and patient, as well as hospital costs.^{5,9}

Considering the need to cover all these aspects, the creation of a protocol that encompasses pre-, peri-, and post-operative measures is a valid strategy to standardize care and increase the safety of the procedure. The present study aimed to propose a multidisciplinary protocol to standardize the care of patients undergoing THA and to evaluate the effectiveness of the protocol after its implantation.

Material and methods

An institutional protocol was created for the management of patients undergoing THA. This protocol is divided into pre-, peri-, and post-operative measures and includes medical, nursing, and physiotherapy professionals.

It begins with the request of preoperative exams in order to identify possible risk factors and infection sites (urinary and airway), and a cardiologist and anesthesiologist evaluation; when necessary, blood components and the intensive care unit bed (ICU) are reserved. The patient is advised to shower

preoperatively using chlorhexidine detergent, and to purchase elastic compression stockings for postoperative use.

After the preoperative procedures are concluded, the patient is admitted on the day of surgery; the recommendations from the World Health Organization (WHO) manual for safe surgery are followed.¹⁰ In the operating room, up to 60 minutes before the incision is made, antibiotic prophylaxis is administered; shaving is performed only if necessary. After surgery, elastic compression stockings are placed, followed by the final control X-ray. Antibiotic prophylaxis is maintained for 24 h, and thromboprophylaxis is initiated with 40 mg enoxaparin daily until discharge.

Once the patient is in the room, the internal medicine staff initiates monitoring. Gait training is initiated on the first postoperative day (POD) by the attending physician, and a blood count is requested. The dressing is changed after 48 h, and the discharge is planned for up to 96 h. At hospital discharge, the patient is advised on the use of the anticoagulant for five weeks after surgery, analgesia according to the pain, gait with a walker and always with an assistant, home physiotherapy, identification of risk signs that require reevaluation, and measures to prevent prosthesis dislocation. This protocol was implemented in May 2012.

The charts of patients who underwent elective THA for hip arthrosis treatment were retrospectively evaluated. A database was created in Microsoft Excel. Initially, information on 100 patients operated in 2011 and 2012 was entered into the database; 50 patients had been operated on before protocol implantation and 50, after. This sample included patients with previous hip surgeries, those with advanced deformities, and those with hip infection sequelae. Three patients who were operated on before implantation of the multidisciplinary protocol were excluded (two were submitted to THA due to femoral neck fracture and the necessary information was not available for the third patient), as well as two who were operated on after protocol implantation (in both cases, surgery was due to fracture). Thus, 95 patients were evaluated, 47 of whom were operated before and 48 after protocol implantation.

The studied variables were: infection, dislocation, DVT, 90-day clinical complications, length of hospital stay, emergency room (ER) visits, re-admission for any reason, and 90-day mortality. Clinical complication was defined as any alteration not

Download English Version:

<https://daneshyari.com/en/article/8600507>

Download Persian Version:

<https://daneshyari.com/article/8600507>

[Daneshyari.com](https://daneshyari.com)