



**Chemotherapy use, end-of-life care, and costs of care among patients diagnosed with Stage****IV pancreatic cancer**

Yuhua Bao, PhD<sup>1,2</sup>, Renee C. Maciejewski, BS<sup>3</sup>, Melissa M. Garrido, PhD<sup>4,5</sup>, Manish A. Shah,  
MD<sup>6</sup>, Paul K. Maciejewski, PhD<sup>3,7-8</sup>, Holly G. Prigerson, PhD<sup>3,8</sup>

<sup>1</sup> Department of Healthcare Policy & Research, Weill Cornell Medical College, New York, New York, U.S.A.

<sup>2</sup> Department of Psychiatry, Weill Cornell Medical College, New York, New York, U.S.A.

<sup>3</sup> Center for Research on End-of-Life Care, Weill Cornell Medical College, New York, New York, U.S.A.

<sup>4</sup> Department of Geriatrics and Palliative Medicine, Mount Sinai School of Medicine, New York, New York, U.S.A.

<sup>5</sup> James J Peters Veteran Affairs Medical Center, Bronx, New York, U.S.A.

<sup>6</sup> Meyer Cancer Center at Weill Cornell Medicine, Medical Oncology/Solid Tumor Program, New York, New York, U.S.A.

<sup>7</sup> Department of Radiology, Weill Cornell Medical College, New York, New York, U.S.A.

<sup>8</sup> Division of Geriatrics and Palliative Medicine, Department of Medicine, Weill Cornell Medical College, New York, New York, U.S.A.

Corresponding Author: Yuhua Bao, 425 East 61<sup>st</sup> Street, New York, New York, U.S.A.

(O) 646 9628037, (F) 646 9620105, [yub2003@med.cornell.edu](mailto:yub2003@med.cornell.edu),

Number of tables: 3

Number of figures: 0

Number of references: 39

Word count: 3,412

Download English Version:

<https://daneshyari.com/en/article/8605571>

Download Persian Version:

<https://daneshyari.com/article/8605571>

[Daneshyari.com](https://daneshyari.com)