

Accepted Manuscript

Additives used to reduce perioperative opioid consumption 1: Alpha2-agonists

Peter H. Tonner, M.D.

PII: S1521-6896(17)30081-2

DOI: [10.1016/j.bpa.2017.10.004](https://doi.org/10.1016/j.bpa.2017.10.004)

Reference: YBEAN 969

To appear in: *Best Practice & Research Clinical Anaesthesiology*


Received Date: 18 January 2017

Revised Date: 25 September 2017

Accepted Date: 26 October 2017

Please cite this article as: Tonner PH, Additives used to reduce perioperative opioid consumption 1: Alpha2-agonists, *Best Practice & Research Clinical Anaesthesiology* (2017), doi: 10.1016/j.bpa.2017.10.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Additives used to reduce perioperative opioid consumption

1: Alpha2-agonists

Peter H. Tonner, M.D.

Dept. of Anaesthesiology and Intensive Care Medicine, Saale-Unstrut Klinikum, Naumburg,
Germany

Correspondence to:

Prof. Dr. Peter H. Tonner

Dept. of Anaesthesiology and Intensive Care Medicine

Saale-Unstrut Klinikum

Humboldtstr. 31

06618 Naumburg

Germany

Tel: +49-3445-722043

Fax: +49-3445-721104

Email: peter.tonner@klinikum-burgendlandkreis.de

Download English Version:

<https://daneshyari.com/en/article/8610884>

Download Persian Version:

<https://daneshyari.com/article/8610884>

[Daneshyari.com](https://daneshyari.com)