Accepted Manuscript

Different protocols used today to achieve total opioid-free general anesthesia without locoregional blocks

Eckhard Mauermann, MD, MSc, Wilhelm Ruppen, MD, Oliver Bandschapp, MD

PII: \$1521-6896(17)30083-6

DOI: 10.1016/j.bpa.2017.11.003

Reference: YBEAN 971

To appear in: Best Practice & Research Clinical Anaesthesiology


Received Date: 17 January 2017

Revised Date: 8 August 2017

Accepted Date: 8 November 2017

Please cite this article as: Mauermann E, Ruppen W, Bandschapp O, Different protocols used today to achieve total opioid-free general anesthesia without locoregional blocks, *Best Practice & Research Clinical Anaesthesiology* (2017), doi: 10.1016/j.bpa.2017.11.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Different protocols used today to achieve total opioid-free

general anesthesia without locoregional blocks

Eckhard Mauermann, MD, MSc^{1,2}, Wilhelm Ruppen, MD¹, Oliver Bandschapp,

MD^{1*}

University Hospital Basel, Department for Anesthesia, Surgical Intensive Care, 1

Prehospital Emergency Medicine and Pain Therapy, Spitalstrasse 21, 4031 Basel,

Switzerland

University Hospital Ghent, Department of Anesthesiology and Perioperative 2

Medicine, Building K12-C 2nd floor, De Pintelaan 185, B-9000 Ghent, Belgium

Corresponding author:

O. Bandschapp

E-mail address: oliver.bandschapp@usb.ch

[Word count: approx. 9200]

Download English Version:

https://daneshyari.com/en/article/8610888

Download Persian Version:

https://daneshyari.com/article/8610888

<u>Daneshyari.com</u>