




# REVISTA BRASILEIRA DE ANESTESIOLOGIA

Publicação Oficial da Sociedade Brasileira de Anestesiologia  
[www.sba.com.br](http://www.sba.com.br)



## SCIENTIFIC ARTICLE

# Functional residual capacity increase during laparoscopic surgery with abdominal wall lift<sup>☆</sup>

Hiroshi Ueda<sup>a</sup>, Takuo Hoshi<sup>b,\*</sup>

<sup>a</sup> Department of Anesthesiology and Critical Care Medicine, Ibaraki Prefectural Central, Ibaraki, Japan

<sup>b</sup> Department of Anesthesiology and Critical Care Medicine, Ibaraki Clinical and Training Center, Tsukuba University Hospital, Ibaraki, Japan

Received 16 November 2015; accepted 4 December 2015

### KEYWORDS

Abdominal wall lift;  
Functional residual capacity;  
Laparoscopic surgery

### Abstract

**Background and objectives:** The number of laparoscopic surgeries performed is increasing every year and in most cases the pneumoperitoneum method is used. One alternative is the abdominal wall lifting method and this study was undertaken to evaluate changes of functional residual capacity during the abdominal wall lift procedure.

**Methods:** From January to April 2013, 20 patients underwent laparoscopic cholecystectomy at a single institution. All patients were anesthetized using propofol, remifentanyl and rocuronium. FRC was measured automatically by Engstrom Carestation before the abdominal wall lift and again 15 min after the start of the procedure.

**Results:** After abdominal wall lift, there was a significant increase in functional residual capacity values (before abdominal wall lift  $1.48 \times 10^3$  mL, after abdominal wall lift  $1.64 \times 10^3$  mL) ( $p < 0.0001$ ). No complications such as desaturation were observed in any patient during this study.

**Conclusions:** Laparoscopic surgery with abdominal wall lift may be appropriate for patients who have risk factors such as obesity and respiratory disease.

© 2016 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### PALAVRAS-CHAVE

Elevador da parede abdominal;  
Capacidade residual funcional;  
Cirurgia laparoscópica

### Aumento da capacidade residual funcional durante cirurgia laparoscópica com elevador da parede abdominal

### Resumo

**Justificativa e objetivos:** O número de cirurgias laparoscópicas realizadas está aumentando a cada ano e, na maioria dos casos, o método pneumoperitônio é o escolhido. Uma alternativa é o método de elevação da parede abdominal, e este estudo foi realizado para avaliar as

<sup>☆</sup> This study was carried out by the Ibaraki Prefectural Central Hospital.

\* Corresponding author.

E-mail: [thoshi@md.tsukuba.ac.jp](mailto:thoshi@md.tsukuba.ac.jp) (T. Hoshi).

<http://dx.doi.org/10.1016/j.bjane.2015.12.003>

0104-0014/© 2016 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

alterações da capacidade residual funcional durante o procedimento de elevação da parede abdominal.

**Métodos:** De janeiro a abril de 2013, 20 pacientes foram submetidos à colecistectomia laparoscópica em uma única instituição. Todos os pacientes foram anestesiados com propofol, remifentanil e rocurônio. A CRF foi medida automaticamente usando o Engström Carestation antes da elevação da parede abdominal e, novamente, 15 minutos após o início do procedimento.

**Resultados:** Após elevar a parede abdominal, um aumento significativo foi observado nos valores da capacidade residual funcional (antes da elevação da parede abdominal:  $1,48 \times 10^3$  mL; após a elevação da parede abdominal:  $1,64 \times 10^3$  mL) ( $p < 0,0001$ ). Não houve complicações, como dessaturação, em nenhum paciente durante este estudo.

**Conclusões:** A cirurgia laparoscópica com elevador da parede abdominal pode ser apropriada para pacientes com fatores de risco como obesidade e doenças respiratórias.

© 2016 Sociedade Brasileira de Anestesiologia. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The use of laparoscopic techniques in surgery is increasing year by year. There are two major techniques for laparoscopic surgery, pneumoperitoneum method and abdominal wall lift method. Establishing a pneumoperitoneum is a major technique for laparoscopic surgery, while lifting procedure is a minor technique. A patient's pulmonary functions are affected by various factors during laparoscopic surgery. Functional residual capacity (FRC) is decreased by the supine position and the induction of anesthesia<sup>1-3</sup> during surgery. The decrease in FRC may cause hypoxemia due to increases in blood flow where gas exchange is not taking place. Furthermore, pulmonary compliance is decreased by the pneumoperitoneum method,<sup>4</sup> but not by the abdominal wall lift.<sup>5</sup>

No previous study has evaluated FRC during laparoscopic surgery with abdominal wall lift. In the current study we test our hypothesis that abdominal wall lift method increases FRC.

## Methods

Our study plan was approved from the institutional review board. We retrospectively studied 20 adult patients who underwent elective laparoscopic cholecystectomy with abdominal wall lift from January to April 2013 at Ibaraki Prefectural Central Hospital. We did not exclude anyone from this study. During the procedure, the right costal arch and the navel area were lifted with a subcutaneous wire using the abdominal wall-lift system (Mizuho Ika, Tokyo, Japan). FRC was measured repeatedly by Engstrom Carestation (GE Health Care, UK Ltd., Buckinghamshire, UK). It takes several minutes in the measurement of FRC. We use average of two to three measurements before the abdominal wall lift and 15–30 min after the start of the procedure.

## The anesthesia and monitoring

The patients were not given any sedative drugs before surgery. General anesthesia was induced with remifentanil

$0.2 \mu\text{g kg}^{-1} \text{min}^{-1}$  and target controlled infusion of propofol (target concentration of plasma was  $3 \mu\text{g mL}^{-1}$ ).

Rocuronium was used for neuromuscular block. Tracheal intubation was performed with tracheal tubes of internal diameters of 7 and 8 mm used for female and male patients, respectively. Anesthesia was maintained with propofol and remifentanil to maintain the bispectral index between 40 and 60 and the systolic pressure at  $\pm 30\%$  of pre-anesthetic values. To maintain neuromuscular block, rocuronium was given intermittently and their train-of-four ratio of 0% was confirmed. The lungs were ventilated mechanically with 30–40% oxygen in air, tidal volume  $8 \text{ mL kg}^{-1}$ , at a respiratory rate of  $10 \text{ min}^{-1}$ .

During anesthesia all patients were monitored by electro cardiogram, non-invasive blood pressure, pulse oximetry, bispectral index, and train-of-four.

## Statistical analysis

Based on a previous Japanese study on FRC changes in anesthetized and intubated patients,<sup>6</sup> power analysis revealed that a minimum sample size of 17 was required in order to detect a difference of 15% in FRC increasing after abdominal wall lift ( $\beta = 0.80$ ,  $\alpha = 0.05$ ). Data are presented as mean ( $\pm$ SD). Comparisons are made between the FRC before and after abdominal wall lift by paired *t*-test (Stat View 5.0, SAS Institute, NC, USA) and  $p < 0.05$  is considered to be a significant difference.

## Results

Patient characteristics are summarized in Table 1. There were 3 obese patients with a Body Mass Index (BMI) of  $> 30 \text{ kg m}^{-2}$ . All patients were included in the statistical analysis. After abdominal wall lift, there was a significant increase of functional residual capacity values (before abdominal wall lift  $1.48 \times 10^3$  mL, after abdominal wall lift  $1.64 \times 10^3$  mL) ( $p < 0.0001$ ).

Linear regression analysis showed that there was a univariate correlation between BMI and increase of FRC (Fig. 1).

Download English Version:

<https://daneshyari.com/en/article/8612149>

Download Persian Version:

<https://daneshyari.com/article/8612149>

[Daneshyari.com](https://daneshyari.com)