



REVISTA BRASILEIRA DE ANESTESIOLOGIA

Official Publication of the Brazilian Society of Anesthesiology
www.sba.com.br



CLINICAL INFORMATION

Neuraxial anesthesia after local anesthesia for management of percutaneous vertebroplasty complication during vertebroplasty

Hüseyin Balkarlı^a, Mesut Kılıç^b, İbrahim Öztürk^{c,*}

^a Akdeniz University Faculty of Medicine, Orthopaedics and Traumatology, Antalya, Turkey

^b Ondokuzmayıs University Faculty of Medicine, Orthopaedics and Traumatology, Samsun, Turkey

^c Dışkapı Yıldırım Beyazıt Education and Research Hospital, Anesthesiology and Reanimation, Ankara, Turkey

Received 9 June 2014; accepted 16 July 2014

KEYWORDS

Neuraxial anesthesia;
Percutaneous
vertebroplasty;
Vertebra;
Fracture

Abstract Percutaneous vertebroplasty is a relatively safe, simple and commonly performed interventional procedure for the management of vertebral compression fractures. However, serious complications are rarely reported in the procedure. Those are pulmonary embolism, severe infection, paraplegia and an occurrence of a new fracture in an adjacent vertebra after vertebroplasty. Acute complications are generally associated with the procedure. We present the case of neuraxial anesthesia, developed after local anesthesia with 8 mL of 2% prilocaine, in a 68-year-old woman who underwent percutaneous vertebroplasty after an osteoporotic collapsed fracture in the L₁ vertebra due to trauma. To our knowledge, this is the first case in the literature.

© 2014 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. All rights reserved.

PALAVRAS-CHAVE

Bloqueio neuraxial;
Vertebroplastia
percutânea;
Vértebra;
Fratura

Bloqueio neuroaxial após anestesia local para manejo de complicação de vertebroplastia percutânea durante vertebroplastia

Resumo Vertebroplastia percutânea é um procedimento intervencionista relativamente seguro, simples e comumente realizado para tratar fraturas por compressão vertebral. No entanto, as complicações graves relacionadas ao procedimento são raramente relatadas, incluindo embolia pulmonar, infecção grave, paraplegia e a ocorrência de uma nova fratura em vértebra adjacente após a vertebroplastia. Complicações agudas são geralmente associadas ao procedimento. Apresentamos o caso de bloqueio neuroaxial, realizado após anestesia local

* Corresponding author.

E-mail: drozturk28@gmail.com (İ. Öztürk).

com 8 mL de prilocaína a 2%, em uma mulher de 68 anos de idade, submetida à vertebroplastia percutânea após fratura osteoporótica na vértebra L₁ devido a trauma. De acordo com nossa pesquisa, este é o primeiro caso na literatura.

© 2014 Sociedade Brasileira de Anestesiologia. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

Percutaneous vertebroplasty (PV) is a relatively safe, simple, and commonly performed procedure for the management of vertebral compression fractures, but serious complications have been reported, including pulmonary embolism,¹ severe infection,² paraplegia³ and an occurrence of a new fracture in an adjacent vertebra after vertebroplasty.⁴ Acute complications are mostly related to surgical procedures, which include leakage of bone cement, infection, and fracture along the path of the vertebroplasty cannula.^{5,6}

We present a rare case of neuraxial anesthesia correlated with local anesthesia during a surgical PV using the transpedicular approach.

Case report

A 68-year-old and 75 kg weighing female patient was admitted to the Emergency Department of Tavas Community Hospital with back pain due to trauma following falling down at home. The patient had an osteoporotic collapsed fracture in the L₁ vertebra, which was treated conservatively with bedrest, corset and analgesic medications. At the 3rd week follow-up examination, pain had not reduced and on the direct radiograph more than 50% of the fractured L₁ vertebra was observed to have collapsed compared to the upper adjacent vertebra (Fig. 1). PV was recommended for the patient.

At the preoperative evaluation of patient, laboratory tests, electrocardiogram and X-ray of lung were normal. There was no chronic systemic disease. And her physical status was evaluated as level 1 according to American Society of Anesthesiologist's scale. The preoperative visual analogue pain scale score (VAS) was 8. After obtaining informed consent, patient's fracture line was located with fluoroscopy in prone position and then under fluoroscopic control, local anesthesia of 8 mL of 2% prilocaína was applied from the pedicle entrance as far as the subcutaneous tissue. No neurological deficit was determined following local anesthesia. After location of the pedicle entrance of the fractured vertebra under fluoroscopy, an 11G needle was entered from the pedicle. When the needle reached 2/3 anterior of the vertebral corpus on the lateral view, the anterior-posterior view was regarded and that the tip of the needle was at the same level as the spinous process. The needle position was checked by fluoroscopy. At this stage, the neurological examination of the patient was normal. After confirmation

of the needle tip location, 8 mL of cement was injected into the fracture line (Fig. 2).

During and following the cementing, the neurological examination of the patient was normal. After setting of the cement, the patient was placed in a supine position. Immediately after the patient was positioned supine, a bilateral lower extremity loss of strength (2/5) and sensation deficit occurred. It was noticed by the operator during neurological examination. Blood pressure was measured as 90/50 mmHg and 500 mL of 0.9% NaCl was administered. As the sensation loss, at T₁₀ level at first examination, reached up to T₅. The patient was moved into the beach-chair position to prevent further development of neuraxial block.

The patient was again evaluated neurologically, and bilateral loss of strength (0/5) in the lower extremities and anesthesia were observed. 6 mg betamethasone was administered intramuscularly. The patient was evaluated with direct radiographs at operating theater and cement was not observed in canal (Fig. 3). 30 min after the onset of the sensation deficit, blood pressure was measured as 110/90 mmHg and the deficit had returned to the level of the umbilicus.

On evaluation of patient by computer tomography, no cement was observed in canal (Fig. 4). At the 4th hour of follow-up sensation and motor deficit was completely resolved. The patient was mobilized. We did not observe hypotension or bradycardia and no inotropic drug was needed.

Discussion

Vertebroplasty has been widely used to treat vertebral body compression fractures (VCFs) caused by varied pathologies including hemangioma, multiple myeloma, osteolytic metastases, and primary or secondary osteoporosis.⁷ VCFs are the most common type of osteoporotic fractures, resulting in severe back pain, spinal deformity, muscle atrophy, prolonged hospitalization, and potential risk of increased mortality.⁸

In meta-analysis of PV, low complication rate of 0.4–3.8% has been reported.⁹ Overall complication rates due to PV are reported to be very low. However, the procedure is not free of risk. Rare, but serious complications have been observed during PV application such as pulmonary embolism, cardiac perforation, epidural cement extravasation paraspinal, intradisc or venous system cement leakage, infection or fractures of adjacent vertebra.

Download English Version:

<https://daneshyari.com/en/article/8612202>

Download Persian Version:

<https://daneshyari.com/article/8612202>

[Daneshyari.com](https://daneshyari.com)