

20th International Congress of Chemical and Process Engineering CHISA 2012
25 – 29 August 2012, Prague, Czech Republic

Biocatalysis in ionic liquid: degradation of phenol by laccase

A. P. M. Tavares a*, B. Pinho, O. Rodriguez, E. A. Macedo

*LSRE - Laboratory of Separation and Reaction Engineering - Associate Laboratory LSRE/LCM, Faculdade de Engenharia,
Universidade do Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal*

Abstract

The aim of this work was to optimize the reaction conditions for the degradation of phenol in aqueous and ionic liquid (IL)-aqueous solution by biocatalysis using two commercial laccases. In order to evaluate the effect of IL on laccase-mediated reactions, the IL 1-ethyl-3-methylimidazolium ethylsulfate, [C₂mim][EtSO₄] was selected as reaction media for this study. Preliminary studies for phenol degradation were carried out in a lab-scale stirred batch reactor with 2 commercial laccases (from *Trametes versicolor*; Sigma and from *Aspergillus*; Novozymes). The synthetic effluent was prepared with phenol in appropriated buffer solutions, with pH varying from 3.0 to 9.0 and different enzyme mediators: 2,2-azinobis(3-ethylbenzothiazoline-6-sulfonate) (ABTS), N-hydroxyacetanilide (NHA) and (2,2,6,6-tetramethylpiperidin-1-yloxy) (TEMPO). For the optimization of reaction conditions, laccase concentration was varied in the range 1000-2000 U/L, pH in the range 6.0-7.0 and IL concentration in the range 10-20% v/v. For comparison, acetonitrile was used as organic solvents at the same conditions. Quantitative analysis of phenols and by-products formed were determined by HPLC (LaChrom Elite HPLC) with a column RP-18 LichroCART at 25°C. In conclusion, laccase (from the class of oxidases) was shown to possess catalytic activity for the degradation of phenol in systems containing ionic liquids. This finding opens up promising perspectives for applying these environmentally benign solvents to a broad range of important oxidative biotransformations.

© 2012 Published by Elsevier Ltd. Selection under responsibility of the Congress Scientific Committee (Petr Kluson) Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Keywords: Laccase; ionic liquid; biocatalytic reaction; phenol

* Corresponding author. Tel.: +351 22 0413606;
E-mail address: atavares@fe.up.pt.

1. Introduction

Industrial processes generate a variety of molecules that may pollute air, soil and water with negative impacts for ecosystems and humans (toxicity, carcinogenic and mutagenic properties). Due to this problem, research has been conducted to investigate the new possibilities of applying enzymes for organic waste treatment. It is known that several types of industrial and agricultural wastes contain phenolic compounds: paints, pesticides, wastes from coal conversion, polymeric resins and petroleum. Due to the fact that the classic treatments of industrial effluents containing phenolic compounds have not yet been able to solve all the associated problems involved or present high costs of process, there is a great need to develop an economical, effective and innovative way for the treatment of these wastes [1]. The potential advantages of the enzymatic treatment when compared to conventional process include: operation at high and low contaminant concentrations; operation over a wide range of pH and temperature. The use of enzymes has been standard in several industries for many years. Only recently they have been studied as a means for enhancing bioremediation. Enzymes possess the ability to break down bonds within organic compounds and/or catalyze their transformation into less toxic and more biodegradable forms. Different studies show that ligninolytic enzymes (laccases) can degrade a wide variety of recalcitrant compounds. Furthermore, the number of laccase substrates can be extended by addition of a redox mediator [2].

Currently, the enzymatic oxidation of phenolic compounds with poor water solubility by enzymatic catalysis is carried out in nonaqueous systems, and the use of organic solvents is required to dissolve them. As disadvantages, the molecules of solvent may directly interact with the enzymes, thus changing their structure, causing irreversible inactivation of the enzymes. Most solvents used present toxicity and low biodegradability. In order to develop a green biotechnological solution, the use of Ionic Liquids (ILs) as solvent or co-solvent is an alternative extremely necessary and interesting [3]. ILs are defined as substances composed only by ions which are liquid at room temperature. Among their properties, publications usually highlight their negligible vapor pressure, thermal and chemical stability, and the possibility to “design” their physicochemical properties by the suitable choice of the anion and the cation [4]. Because of their negligible vapor pressure, ILs are claimed as “green” alternatives for volatile organic compounds (VOCs). Moreover, several ILs have demonstrated to be adequate solvents for enzyme reactions [5].

2. Materials and Methods

2.1 Chemicals

Laccases from *Trametes versicolor* and from *Aspergillus* were purchased from Sigma and kindly provided by Novozymes (Denmark), respectively. Phenol 99% was purchased from Merck. 2,2-azinobis(3-ethylbenzothiazoline-6-sulfonate) (ABTS), N-hydroxyacetanilide (NHA) or (2,2,6,6-tetramethylpiperidin-1-yloxy) (TEMPO) were obtained from Sigma-Aldrich. The IL 1-ethyl-3-methylimidazolium ethylsulfate, [C₂mim][EtSO₄] was synthesized in our laboratory, following a procedure available in the literature [6].

Download English Version:

<https://daneshyari.com/en/article/861578>

Download Persian Version:

<https://daneshyari.com/article/861578>

[Daneshyari.com](https://daneshyari.com)