



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Case report

The confusion between asthma and subglottic stenosis can cause an adverse event during intubation. A case report[☆]



Gilberto Marrugo-Pardo^{a,b,*}, Maria Camila Villalobos-Aguirre^a

^a ENT Department, School of Medicine, Universidad Nacional de Colombia, Bogotá, Colombia

^b Fundación Hospital de la Misericordia, Bogotá, Colombia

ARTICLE INFO

Article history:

Received 13 December 2016

Accepted 21 July 2017

Available online 22 November 2017

Keywords:

Asthma

Spirometry

Laryngostenosis

Respiratory sounds

Case Report

ABSTRACT

Wheezing and stridor are a frequent reason for consultation, as well as the diagnosis of asthma. Any cases of asthma with failed clinical response to treatment should undergo additional testing leading to differential diagnoses. The case herein presented emphasizes the importance of having a knowledgeable surgical team that interprets lung function tests correctly.

This is a case of a 13-year-old girl who underwent corrective surgery of a right tympanic perforation, with a history of difficult to control exercise-induced asthma. At the time of intubation, Cotton's grade III laryngeal stenosis was identified that required intervention prior to the scheduled procedure. Emphasis shall be placed on the importance of interdisciplinary work when evaluating these patients.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

La confusión entre asma y estenosis subglótica puede causar un evento adverso en la intubación. Reporte de un caso

RESUMEN

Las sibilancias y el estridor son un motivo de consulta frecuente, así como el diagnóstico de asma. Aquellos casos de asma sin respuesta clínica al tratamiento deben ser estudiados con pruebas adicionales para realizar diagnósticos diferenciales. Presentamos un caso clínico que resalta la importancia de que todo el equipo quirúrgico conozca e interprete correctamente las pruebas de función pulmonar.

Presentamos el caso de una adolescente de 13 años, llevada a cirugía para corrección de una perforación timpánica derecha, con antecedente de asma inducida por ejercicio de difícil control. Al momento de la intubación se encontró una estenosis laríngea grado III de

Palabras clave:

Asma

Espirometría

Laringoestenosis

Ruidos respiratorios

Informe de caso

[☆] Please cite this article as: Marrugo-Pardo G, Villalobos-Aguirre MC. La confusión entre asma y estenosis subglótica puede causar un evento adverso en la intubación. Reporte de un caso. Rev Colomb Anestesiolog. 2017;45:82-85.

* Corresponding author at: ENT Department, School of Medicine, Universidad Nacional de Colombia, Carrera 45 #26-85, Bogotá, Colombia. E-mail address: gemarrugop@unal.edu.co (G. Marrugo-Pardo).

2256-2087/© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Cotton que fue necesario intervenir antes de realizar la cirugía programada. Resaltamos la importancia del trabajo interdisciplinario en la evaluación de estos pacientes.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Wheezing is described as a constant noise lasting more than 0.25 s that is caused by the oscillation of the opposing airway walls. It may occur during inspiration or expiration and may be caused by narrowing of the airway. Stridor is a high-frequency sound caused by air turbulence inside the upper airway. Though wheezing and stridor are clinically different, they are often mistaken.

One of the causes of stridor is laryngotracheal stenosis resulting from an abnormal narrowing along the glottis and the carina. Before 1960 the condition was a congenital etiology, but with the wide use of orotracheal intubation, its incidence rose from 0.9%¹ to 24.5%.² The most frequent benign cause is tracheal intubation and the most frequent malignant cause is squamous cell carcinoma. The most widely accepted stenosis-associated factor following endotracheal intubation is the duration of the intubation. Notwithstanding the existing evidence about its relevance, the occurrence of cases after short endotracheal intubations suggests that laryngeal lesions are multifactorial, involving patient as well as intubation-associated factors.³

Furthermore, wheezing secondary to exercise-induced bronchoconstriction is the primary indicator of asthma exacerbations and the last to be resolved.⁴ In case of an asthmatic patient, a differential diagnosis shall be done between inspiratory symptoms during exercise, typically occurring at the end of the exercise session, and exercise-induced asthma, with onset of symptoms between 3 and 15 min after exercise. Finally, the possibility of exercise-induced laryngeal obstruction, associated to adduction of the vocal folds as a secondary phenomenon shall also be kept in mind.⁵

When studying asthma, the flow-volume curve evidences highly suggestive changes, although it is a rare cause of dyspnea and wheezing during exertion. However, without a clinical suspicion, the symptoms may be confused for asthma. Delays in making the diagnosis may result in increased morbidity.⁶

Lung function tests are useful for diagnosing unexplained respiratory symptoms and for monitoring patients with known respiratory disease. Generally speaking, patients over 5 years of age are considered able to cooperate with the proper performance of the procedure. A FVC ratio <70% in adults and <80% in children suggests obstructive pathology and the bronchodilator response must be evaluated. If the FVC is more than 12% between 5 and 18 years of age, or more than 12% and >200 ml in adults, a reversible obstruction suggestive of asthma may be considered. However, if these premises are not met, an irreversible obstruction should be suspected and differential diagnoses must be evaluated.⁷ When the flow-volume curve is flat, the most frequent cause is submaximal effort, but then a central or upper airway obstruction must be considered.^{8,9}

The Empey¹⁰ index has been classically used to assess airway obstruction using spirometry. It measures the ratio between VEF1 (ml) and PEFr (l/min); if the ratio is over 10, this is considered evidence of laryngotracheal stenosis.

Properly diagnosing upper or lower respiratory disease is vitally important in the pre-surgical setting, to prevent unexpected difficult airway cases and getting the surgical team ready to optimize patient outcomes.

Clinical case

This case is a 13-year old female patient with a history of patent ductus arteriosus surgically corrected 7 days after birth. The patient did not require orotracheal intubation following the procedure. Additionally, the patient has a history of asthma and is managed with inhaled steroids, with persistent symptoms when she exercises and follow-up by pulmonology. She came to the ENT department of the Fundación Hospital de la Misericordia because of hearing loss and repeated episodes of otitis media during her childhood. A sub-total right tympanic perforation was identified, and the patient was scheduled for type 1 tympanoplasty as an ambulatory procedure. During the pre-anesthesia evaluation, the patient was classified as ASA II because of her history of asthma and continued management was recommended. Following the anesthetic induction, difficulty for intubation with a #6.0 cuffed endotracheal tube was reported, with evidence of subglottic concentric stenosis during laryngoscopy; the anesthetist then switched to a #4.0 cuffed tube. Since the patient was admitted to a high-complexity hospital, she simultaneously underwent laryngeal microendoscopy and bronchoscopy, evidencing a Cotton-Myer III¹¹ concentric subglottic stenosis affecting 70% of the subglottic lumen (Fig. 1), that required laryngeal dilatation and mitomycin and triamcinolone infiltration as per the institutional protocol (Fig. 2). The pre-surgical spirometry was studied, evidencing a curve with discrete flattening of the expiratory peak, reporting all parameters within the normal limits, except for a 60% PEF, with poor response to beta2-agonists (Fig. 3).

Discussion

The British Thoracic Society¹² recommends that unless an alternate diagnosis is more likely, patients with symptoms suggestive of asthma shall be initiated on medical management and further testing is advised if the response is poor.

When clinically evaluating the differential diagnoses, the upper airway obstruction usually involves stridor, contrary to asthma that presents with wheezing. However, there may be

Download English Version:

<https://daneshyari.com/en/article/8616378>

Download Persian Version:

<https://daneshyari.com/article/8616378>

[Daneshyari.com](https://daneshyari.com)