



Research article

Proposal of a score to detect the need for postoperative intensive care unit admission after bariatric surgery



Walid H. Nofal*, Akram M. Amer, Walid A. Mansour

Department of Anaesthesiology, Intensive Care and Pain Management, Faculty of Medicine, Ain-Shams University, Abbasia, Cairo, Egypt

ARTICLE INFO

Article history:

Received 16 June 2017

Revised 30 July 2017

Accepted 16 August 2017

Available online 24 August 2017

Keywords:

Bariatric surgery

Postoperative ICU admission

Score

ABSTRACT

Background: We developed a multi-dimensional score which may help in predicting those patients, undergoing bariatric surgery, who may be in need for postoperative ICU admission and which may also help in avoiding unnecessary admission to the critical care units after bariatric surgery.

Methods: We collected the data of 111 patients who underwent either laparoscopic gastric sleeve or bypass and studied the association between some risk factors related to obesity and their postoperative ICU admission. Those factors found to be statistically significant are included in the final score. The cutoff value of our scoring system is determined by running a Receiver Operating Curve (ROC) analysis.

Results: Forty patients (36%) were admitted to the ICU postoperatively. Our final score includes 7 independent variables; 6 found to be significantly related to post-bariatric surgery ICU admission; these are age, gender, BMI, ASA, obstructive sleep apnea and spirometry results, and the seventh is the history of venous thrombo-embolism. According to the ROC curve analysis, we set the score value of 10 as our cut-off value for the need of postoperative ICU admission. The score median value is 9. Males' odds to be admitted to the ICU after bariatric surgery are 11.9 times higher than females. Also, those with BMI above 50 kg m^{-2} have odds of 29.8 times higher than those below 50 kg m^{-2} .

Conclusions: We propose a scoring system for risk stratification, in which some of the well-known predictor risk factors are included in a simple way to help identify those high-risk patients undergoing bariatric surgery.

Conclusions: Trial registry number: NCT02976649.

© 2017 Publishing services by Elsevier B.V. on behalf of Egyptian Society of Anesthesiologists. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Obesity is a current global threat, with an estimated 1.9 billion over weight adults, plus more than 600 million are suffering obesity [1]. Those with body mass index (BMI) of 30 kg m^{-2} or more constitute 27.7% of adult population in Egypt [2]. In parallel; there is an exponential increase of bariatric surgeries. Male sex, above 50 years, BMI more than 60 kg m^{-2} , cardiovascular disease, obstructive sleep apnoea syndrome, venous stasis, and intra-operative complications are known risk factors for intensive care unit (ICU) admission after bariatric surgery [3].

Nowadays, the prediction of ICU need post bariatric surgery depends on diverse protocols ranging from being BMI based to co-morbidities based overlooking in most the physiological reserve of such category of patients. However, despite the morbidity and

mortality risks, most of patients admitted to ICU did not benefit from its services resulting in inefficient utilization [4].

The aim of this study is to create a multi-dimensional predictive score for the need of post-bariatric surgery ICU admission. First, to have an objective tool for assessment of our patients. Second, incorporating all the risk factors and physiological reserve in one comprehensive score. Third, easy tailoring of the predictive ICU needs of post bariatric surgeries with accumulated data and experiences.

2. Material and methods

This is an observational retrospective study which was approved by the Ain Shams University, Faculty of Medicine Research Ethics Committee (Ref: FMASU R 19/2016) and registered with Clinical trials.gov (ref: NCT02976649). The requirement for written informed consent was waived by the ethics committee.

The first step in our work was to develop a preliminary score. This score includes independent factors we considered as risk fac-

Peer review under responsibility of Egyptian Society of Anesthesiologists.

* Corresponding author.

E-mail addresses: drwalid_nofal@med.asu.edu.eg, wahid_hamed_nofal@yahoo.com (W.H. Nofal).

tors for the need for postoperative ICU admission after bariatric surgery. We applied this score on pilot group of patients; first to find those factors which are significantly related to the need for post-bariatric surgery ICU admission, so that to be included in the final score and second to determine the cutoff score value for the prediction of our dependent factor; the need for ICU admission. We collected the data of 111 patients who have been operated upon for either laparoscopic gastric sleeve or laparoscopic gastric bypass during the period from January to June 2016. All the surgeries were performed in Ain Shams University hospitals during the previously mentioned period. The data were collected from the database registry of the general surgery department in our University Hospital. Patients were indicated for surgery if their body mass index (BMI) $\geq 35 \text{ kg m}^{-2}$ with obesity-related comorbidities or $\geq 40 \text{ kg m}^{-2}$ with or without co-morbidities [5]. Patients were excluded in case of postoperative surgical complications, conversion to open surgery or in case of redo. Unfortunately, the available postoperative data for those admitted to the ICU were only about the duration of mechanical ventilation and the duration of their ICU stay.

The items included in the preliminary score are age, gender, BMI, physical status of the patient according to the American Society of Anaesthesiologist (ASA) classification [6], the presence of obstructive sleep apnoea (OSA). Also, we included the spirometry results as an indicator of pulmonary function, history of venous thrombo-embolic events (VTE), and finally type and duration of surgery. All these data besides the admission to the ICU postoperatively were collected for each patient. The association between each of these items and the admission to ICU postoperatively were examined using Chi-square or Fischer exact test as appropriate. For this univariate analysis, the patients were divided into 2 age groups; below 40 and 40 years old or above. Also, the independent variable BMI was examined by dividing the patients into 2 groups; those with BMI below 50 kg m^{-2} or 50 kg m^{-2} and above.

According to the spirometry results, patients are divided into two groups as either normal or with obstructive or restrictive lung disease.

Also, for the univariate analysis, the patients are classified as ASA II or above II and again as with or without obstructive sleep apnoea.

The diagnosis of co-morbid conditions followed standard recommendations. As an example, Diabetes mellitus required an elevated fasting blood sugar of $\geq 120 \text{ mg/dl}$ on two or more tests on different days and the recommendation of a diabetic diet by the primary care physician, oral hypoglycaemic medications or insulin. Hypertension required sitting repeated blood pressure measurements of $\geq 150 \text{ mmHg}$ systolic and/or $\geq 90 \text{ mmHg}$ diastolic or the use of anti-hypertensive medications. Obstructive Sleep Apnoea required polysomnographic findings demonstrating a respiratory disturbance index ≥ 10 hypopnoeic or apnoeic episodes/hour of sleep.

Per the results of the univariate analysis, only the items with significant association with our dependent factor, the postoperative ICU admission, will be included in the final score.

Finally, we obtained a score sheet of 7 elements found to be significantly related to the need for postoperative ICU admission: Age, Gender, BMI, American society of anaesthesiologist physical status, obstructive sleep apnoea, spirometry results and history of venous thrombo-embolism.

In our final score sheet, age and BMI were further divided into 4 sub groups. Each element will be given a score from 1 to 4 except the gender category which has only 2 possible outcomes will be given a score of either 1 or 2. Male gender was given a higher score because of the known comorbidities related to the android type of obesity. Also, per the ASA classification [6], none of our patients belongs to ASA I physical status (all have BMI above 30) so we started our score sheet with ASA II patients who will be given a

score of 1 and so on. The results of the spirometry; either normal, mild, moderate or severe obstructive or restrictive are given scores from 1 to 4 respectively. The pulmonary functions were evaluated by following the American Thoracic Society (ATS) grades for diagnosing the pattern and quantifying the severity of the defect [7].

The presence of OSA and the degree of venous thrombo-embolic disease will be scored as shown in Table 3.

Thus, the score value of any patient will range from 7 to 25. The sum of all elements was calculated for each patient.

We run a Receiver Operating Characteristic (ROC) curve analysis to determine the score cutoff value that best discriminates between patients who will be in need for postoperative ICU admission and those who will not. This will be the value with the best combination between sensitivity and false positive rate.

Finally, we conducted a binomial logistic regression to ascertain the effects of gender, age, BMI, presence of OSA and results of the spirometry on the likelihood that patients will need post-bariatric surgery ICU admission.

2.1. Statistical analysis

Data were analysed using Statistical Package for Social Science (SPSS) version 21.0. Chicago, Illinois, USA. Quantitative data were expressed as mean \pm standard deviation (SD) or median (range). Qualitative data were expressed as frequency and percentage. Chi-square (X²) and Fischer exact tests were used to compare proportions between two qualitative parameters. P-value < 0.05 was considered significant and P-value < 0.001 was considered highly significant. Receiver Operating Characteristic (ROC) curve was used to evaluate our score and to determine the cut off score value that best discriminates between patients who will need postoperative ICU admission and those who do not. Sample size was determined based on a range of ICU admission after bariatric surgery from 5 to 24% with α error of 0.05 and power of the study 80%.

3. Results

The demographic characteristics of the patients are represented in Table 1; males represent 17.1% of the patients. The highest BMI was 74 while the lowest was 35. Forty patients were admitted to the ICU postoperatively representing about 36% of patients. Of these admitted patients, 31 were admitted because of BMI above 50 with or without abnormal spirometry, the remaining nine were admitted because of abnormal spirometry results or their complaint of OSA or combination of both. Those nine patients who needed mechanical ventilation were weaned in the same day of admission to the ICU. All the patients were discharged the next morning except those who needed mechanical ventilation were discharged after 48 hours. No CPAP masks were applied because of its possible hazardous effect on the anastomosis.

Table 1

Demographic characteristics of the patients and surgeries. OSA, obstructive sleep apnoea; ASA, American society of anaesthesiologists; VTE, venous thromboembolism.

Age (years)	33.44 \pm 9.2 ^a
Male gender	19 (17.1) ^b
BMI (kg m^{-2})	48.78 \pm 8.2 ^a
Patients with OSA	26 (23.4) ^b
ASA physical status III or IV	43 (38.7) ^b
Patients with abnormal spirometry results	41 (36.9) ^b
Patients with history of VTE	4 (3.6) ^b
Type of surgery (Gastric Bypass)	39 (35.1) ^b
Duration of surgery (hours)	2.2 \pm 0.66 ^a
ICU admission (yes)	40 (36) ^b
Postoperative ventilation (yes)	9 (8.1) ^b

^a Data presented as mean \pm SD.

^b Data presented as count (percentage).

Download English Version:

<https://daneshyari.com/en/article/8616672>

Download Persian Version:

<https://daneshyari.com/article/8616672>

[Daneshyari.com](https://daneshyari.com)