

BMD-based assessment of local porosity in human femoral cortical bone

Gianluca Iori, Frans Heyer, Vantte Kilappa, Caroline Wyers, Peter Varga, Johannes Schneider, Melanie Gräsel, Robert Wendlandt, Reinhard Barkmann, Joop van den Bergh, Kay Raum

PII: S8756-3282(18)30218-7
DOI: doi:10.1016/j.bone.2018.05.028
Reference: BON 11664

To appear in: *Bone*

Received date: 13 December 2017
Revised date: 9 May 2018
Accepted date: 25 May 2018

Please cite this article as: Gianluca Iori, Frans Heyer, Vantte Kilappa, Caroline Wyers, Peter Varga, Johannes Schneider, Melanie Gräsel, Robert Wendlandt, Reinhard Barkmann, Joop van den Bergh, Kay Raum , BMD-based assessment of local porosity in human femoral cortical bone. Bon (2017), doi:10.1016/j.bone.2018.05.028

BMD-based assessment of local porosity in human femoral cortical bone

Gianluca Iori[(1)], Frans Heyer[(2,3)], Vantte Kilappa[(4)], Caroline Wyers[(2,3)], Peter Varga[(5)], Johannes Schneider[(1)], Melanie Gräsel[(6)], Robert Wendlandt[(7)], Reinhard Barkmann[(6)], Joop van den Bergh[(2,3)] and Kay Raum[(1)]

1. Berlin-Brandenburg Center for Regenerative Therapies, Charité - Universitätsmedizin Berlin, Germany;
2. Department of Internal Medicine, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Center, Maastricht, The Netherlands;
3. Department of Internal Medicine, VieCuri Medical Center, Venlo, The Netherlands;
4. Mango Solutions, Jyväskylä, Finland;
5. AO Research Institute Davos, Davos, Switzerland;
6. Sektion Biomedizinische Bildgebung, Klinik für Radiologie, Universitätsklinikum Schleswig-Holstein, Campus Kiel, Germany;
7. Universitätsklinikum Schleswig-Holstein, Lübeck, Germany.

Corresponding author: Kay Raum (kay.raum@charite.de)