

Accepted Manuscript

Invited review

Association of diabetes mellitus with prostate cancer grade and prostate-specific antigen in Chinese biopsy population

Ming Chen, Yili Luo, Shaoling Yang, Lu Xu, Nan Li, Hong Li, Shen Qu

PII: S0168-8227(18)30351-6

DOI: <https://doi.org/10.1016/j.diabres.2018.04.022>

Reference: DIAB 7337

To appear in: *Diabetes Research and Clinical Practice*


Received Date: 3 March 2018

Revised Date: 7 April 2018

Accepted Date: 10 April 2018

Please cite this article as: M. Chen, Y. Luo, S. Yang, L. Xu, N. Li, H. Li, S. Qu, Association of diabetes mellitus with prostate cancer grade and prostate-specific antigen in Chinese biopsy population, *Diabetes Research and Clinical Practice* (2018), doi: <https://doi.org/10.1016/j.diabres.2018.04.022>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Association of diabetes mellitus with prostate cancer grade and prostate-specific antigen in Chinese biopsy population

Ming Chen¹, Yili Luo¹, Shaoling Yang¹, Lu Xu¹, Nan Li¹, Hong Li^{1*}, Shen Qu¹

¹ Department of Endocrinology and Metabolism, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai 200072, China;

*Correspondence: Hong Li.

Department of Endocrinology and Metabolism, Shanghai Tenth People's Hospital, Tongji University School of Medicine, No. 301, Middle Yan chang Road, Shanghai 200072, China.

E-mail address: lihong_endo@tongji.edu.cn.

Fax: +86 21 66301004.

Highlights:

- Diabetes was associated with a higher risk of PCa detection.
- As diabetic duration and HbA1c increase, high-grade PCa risk was increased.
- Diabetics had a 22.8% lower geometric mean PSA level than non-diabetics.
- Low PSA levels in diabetics might delay and/or interfere with PCa detection.

Disclosure statement: The authors have no conflict of interest.

Download English Version:

<https://daneshyari.com/en/article/8629827>

Download Persian Version:

<https://daneshyari.com/article/8629827>

[Daneshyari.com](https://daneshyari.com)