

Fasting serum amino acids concentration is associated with insulin resistance and pro-inflammatory cytokines

Sang-Guk Lee, Ye Seal Yim, Yong-ho Lee, Byung-Wan Lee, Hyon-Suk Kim, Kyung-Sup Kim, Yong-Wha Lee, Jeong-Ho Kim

PII: S0168-8227(18)30100-1
DOI: https://doi.org/10.1016/j.diabres.2018.03.028
Reference: DIAB 7285

To appear in: *Diabetes Research and Clinical Practice*

Received Date: 17 January 2018
Revised Date: 10 March 2018
Accepted Date: 20 March 2018

Please cite this article as: S-G. Lee, Y. Seal Yim, Y-h. Lee, B-W. Lee, H-S. Kim, K-S. Kim, Y-W. Lee, J-H. Kim, Fasting serum amino acids concentration is associated with insulin resistance and pro-inflammatory cytokines, *Diabetes Research and Clinical Practice* (2018), doi: https://doi.org/10.1016/j.diabres.2018.03.028

Fasting serum amino acids concentration is associated with insulin resistance and pro-inflammatory cytokines

Sang-Guk Lee [a], Ye Seal Yim [a], Yong-ho Lee [b], Byung-Wan Lee [b], Hyon-Suk Kim [a], Kyung-Sup Kim [c], Yong-Wha Lee [d], Jeong-Ho Kim [a, *]

[a] Departments of Laboratory Medicine, Severance Hospital, Yonsei University College of Medicine, Seoul, Korea

[b] Division of Endocrinology and Metabolism, Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Korea

[c] Department of Biochemistry and Molecular Biology, Yonsei University College of Medicine, Seoul, Korea

[d] Department of Laboratory Medicine and Genetics, Soonchunhyang University Bucheon Hospital, Soonchunhyang University College of Medicine, Bucheon, Korea

*__Corresponding author__: Jeong-Ho Kim M.D., Ph.D.

Department of Laboratory Medicine, Yonsei University College of Medicine

50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea

E-mail: jeongho@yuhs.ac; Tel.: +82-2-2228-2448; Fax: +82-2-313-0956.

Short title: Alteration of amino acid profile in diabetes

Key words: amino acids profile, inflammatory cytokines, insulin resistance, type 2 diabetes mellitus, proteolysis