Accepted Manuscript

Serum cystatin C level is associated with carotid arterial wall elasticity in subjects with type 2 diabetes mellitus: a potential marker of early-stage atherosclerosis

Rei Kaneko, Shojiro Sawada, Ai Tokita, Rieko Honkura, Noriko Tumura, Shinjiro Kodama, Tomohito Izumi, Kei Takahashi, Kenji Uno, Junta Imai, Tetsuya Yamada, Yukiya Miyachi, Hideyuki Hasegawa, Hiroshi Kanai, Yasushi Ishgaki, Hideki Katagiri


PII: S0168-8227(17)30744-1

DOI: https://doi.org/10.1016/j.diabres.2018.02.003

Reference: DIAB 7216

To appear in: Diabetes Research and Clinical Practice

Received Date: 1 August 2017 Revised Date: 29 January 2018 Accepted Date: 1 February 2018



Please cite this article as: R. Kaneko, S. Sawada, A. Tokita, R. Honkura, N. Tumura, S. Kodama, T. Izumi, K. Takahashi, K. Uno, J. Imai, T. Yamada, Y. Miyachi, H. Hasegawa, H. Kanai, Y. Ishgaki, H. Katagiri, Serum cystatin C level is associated with carotid arterial wall elasticity in subjects with type 2 diabetes mellitus: a potential marker of early-stage atherosclerosis, *Diabetes Research and Clinical Practice* (2018), doi: https://doi.org/10.1016/j.diabres.2018.02.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Serum cystatin C level is associated with carotid arterial wall elasticity in subjects with type 2 diabetes mellitus: a potential marker of early-stage atherosclerosis

Rei Kaneko¹, Shojiro Sawada¹, Ai Tokita¹, Rieko Honkura¹, Noriko Tumura¹, Shinjiro Kodama¹, Tomohito Izumi¹, Kei Takahashi¹, Kenji Uno¹, Junta Imai, Tetsuya Yamada¹, Yukiya Miyachi², Hideyuki Hasegawa³, Hiroshi Kanai⁴, Yasushi Ishgaki⁵, Hideki Katagiri¹

- ¹ Department of Metabolism and Diabetes, Tohoku University Graduate School of Medicine, Sendai, Japan
- ² Medical Systems Research and Development Center R&D Management Headquarters, FUJIFILM Corporation, Kaisei, Japan
- ³ Graduate School of Science and Engineering, University of Toyama, Toyama, Japan
- ⁴ Department of Electronic Engineering, Tohoku University Graduate School of Medicine, Sendai, Japan
- ⁵ Department of Internal Medicine, Division of Diabetes and Endocrinology, Iwate Medical University Hospital, Morioka, Japan

Address correspondence to:

Shojiro Sawada, M.D., Ph.D. and Hideki Katagiri, M.D., Ph.D.

Department of Metabolism and Diabetes, Tohoku University Graduate School of Medicine, 2-1 Seiryo-machi, Aoba-ku, Sendai 980-8575,Japan

Phone & Fax: +81-22-717-8228,

E-mail: ssawada@med.tohoku.ac.jp, katagiri@med.tohoku.ac.jp

Download English Version:

https://daneshyari.com/en/article/8629957

Download Persian Version:

https://daneshyari.com/article/8629957

<u>Daneshyari.com</u>