## Accepted Manuscript

Plasma triglycerides as a risk factor for chronic kidney disease in type 2 diabetes mellitus: Evidence from the northeastern of Thailand

Sojib Bin Zaman, Mohd Anisul Karim, Naznin Hossain, Gulam Muhammed Al Kibria, Sheikh Mohammed Shariful Islam


PII: S0168-8227(17)31025-2

DOI: https://doi.org/10.1016/j.diabres.2018.02.011

Reference: DIAB 7224

To appear in: Diabetes Research and Clinical Practice

Received Date: 22 June 2017
Revised Date: 25 December 2017
Accepted Date: 6 February 2018



Please cite this article as: S. Bin Zaman, M. Anisul Karim, N. Hossain, G. Muhammed Al Kibria, S. Mohammed Shariful Islam, Plasma triglycerides as a risk factor for chronic kidney disease in type 2 diabetes mellitus: Evidence from the northeastern of Thailand, *Diabetes Research and Clinical Practice* (2018), doi: https://doi.org/10.1016/j.diabres.2018.02.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **ACCEPTED MANUSCRIPT**

### Title page

Title: Plasma triglycerides as a risk factor for chronic kidney disease in type 2 diabetes

mellitus: Evidence from the northeastern of Thailand

#### **Authors:**

Sojib Bin Zaman<sup>1,2,3</sup>

Mohd Anisul Karim<sup>4</sup>

Naznin Hossain<sup>5</sup>

Gulam Muhammed Al Kibria<sup>6</sup>

Sheikh Mohammed Shariful Islam<sup>7,8</sup>

#### **Affiliations:**

#### **Corresponding author:**

Sojib Bin Zaman; Maternal and Child Health Division, International Centre for Diarrhoeal

Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh

Email: sojibbz@gmail.com

Type of Contribution: Research Paper

Running title: Plasma triglyceride as a risk factor for CKD in T2DM patients

<sup>&</sup>lt;sup>1</sup>Faculty of Public Health, Khon Kaen University, Thailand

<sup>&</sup>lt;sup>2</sup>Institute of Tropical Medicine and International Health, Charite- University Medicine Berlin, Germany

<sup>&</sup>lt;sup>3</sup>Maternal and Child Health Division, International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh

<sup>&</sup>lt;sup>4</sup>Nuffield Department of Population Health, University of Oxford, UK

<sup>&</sup>lt;sup>5</sup>Department of Pharmacology, Dhaka Medical College, Bangladesh

<sup>&</sup>lt;sup>6</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

<sup>&</sup>lt;sup>7</sup>Sydney Medical School, University of Sydney, Australia

<sup>&</sup>lt;sup>8</sup>Cardiovascular Division, The George Institute for Global Health, Australia

#### Download English Version:

# https://daneshyari.com/en/article/8630066

Download Persian Version:

https://daneshyari.com/article/8630066

<u>Daneshyari.com</u>