

Murine remote preconditioning increases glucose uptake and suppresses gluconeogenesis in hepatocytes via a brain-liver neurocircuit, leading to counteracting glucose intolerance

Atsushi Kurabayashi, Chiharu Tanaka, Waka Matsumoto, Seiji Naganuma, Mutsuo Furihata, Keiji Inoue, Yoshihiko Kakinuma

PII: S0168-8227(17)31400-6
DOI: https://doi.org/10.1016/j.diabres.2018.03.009
Reference: DIAB 7266

To appear in: *Diabetes Research and Clinical Practice*

Received Date: 28 August 2017
Revised Date: 21 January 2018
Accepted Date: 2 March 2018

Please cite this article as: A. Kurabayashi, C. Tanaka, W. Matsumoto, S. Naganuma, M. Furihata, K. Inoue, Y. Kakinuma, Murine remote preconditioning increases glucose uptake and suppresses gluconeogenesis in hepatocytes via a brain-liver neurocircuit, leading to counteracting glucose intolerance, *Diabetes Research and Clinical Practice* (2018), doi: https://doi.org/10.1016/j.diabres.2018.03.009

Murine remote preconditioning increases glucose uptake and suppresses gluconeogenesis in hepatocytes via a brain-liver neurocircuit, leading to counteracting glucose intolerance

Atsushi Kurabayashi[a], Chiharu Tanaka[a], Waka Matsumoto[a], Seiji Naganuma[a], Mutsuo Furihata[a], Keiji Inoue[b], Yoshihiko Kakinuma[c]

Department of [a]Pathology and [b]Urology, Kochi Medical School, Kochi, 783-8505, Japan; [c]Department of Physiology, Nippon Medical School Graduate School of Medicine, Tokyo, 113-8602, Japan

Correspondence and requests for materials should addressed to

Yoshihiko Kakinuma M.D., Ph.D.

Graduate School Professor, Department of Physiology, Nippon Medical School Graduate School of Medicine 1－1－5 Sendagi, Bunkyo-ku, Tokyo, 113-8602,

Japan

TEL: +81-3-3822-2131 Ext. 5244; FAX: +81-3-3822-0766

E-mail: k12417853@nms.ac.jp