

Tissue distribution and biochemical characteristics of receptors for sinus gland peptide VII as a crustacean hyperglycemic hormone and vitellogenesis-inhibiting hormone of the kuruma prawn, *Marsupenaeus japonicus*

Chiaki Nagai-Okatani, Shinji Nagata, Hiromichi Nagasawa

PII: S0016-6480(18)30036-4
DOI: https://doi.org/10.1016/j.ygcen.2018.05.010
Reference: YGCEN 12936

To appear in: *General and Comparative Endocrinology*

Received Date: 10 January 2018
Revised Date: 7 May 2018
Accepted Date: 7 May 2018

Please cite this article as: Nagai-Okatani, C., Nagata, S., Nagasawa, H., Tissue distribution and biochemical characteristics of receptors for sinus gland peptide VII as a crustacean hyperglycemic hormone and vitellogenesis-inhibiting hormone of the kuruma prawn, *Marsupenaeus japonicus*, *General and Comparative Endocrinology* (2018), doi: https://doi.org/10.1016/j.ygcen.2018.05.010

Tissue distribution and biochemical characteristics of receptors for sinus gland peptide VII as a crustacean hyperglycemic hormone and vitellogenesis-inhibiting hormone of the kuruma prawn, *Marsupenaeus japonicus*

Chiaki Nagai-Okatani[1], Shinji Nagata[2] and Hiromichi Nagasawa[*]

Department of Applied Biological Chemistry, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Yayoi 1-1-1, Bunkyo-ku, Tokyo 113-8657, Japan.

* Corresponding author.

Hiromichi Nagasawa, Ph.D.

Department of Applied Biological Chemistry, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan.

Tel.: +81-3-5841-5132

Fax.: +81-3-5841-8022

E-mail: anagahi@mail.ecc.u-tokyo.ac.jp