## Accepted Manuscript

Effect of infusion speed of 7.5% hypertonic saline on brain edema in patients with craniocerebral injury: An experimental study

Zhenzhen Jiang, Hongmei Xu, Meilin Wang, Zefu Li, Xinyang Su, Xiaoli Li, Zhenzhu Li, Xuexin Han

PII: S0378-1119(18)30482-7

DOI: doi:10.1016/j.gene.2018.05.005

Reference: GENE 42816

To appear in: Gene

Received date: 4 March 2018 Revised date: 27 April 2018 Accepted date: 2 May 2018

Please cite this article as: Zhenzhen Jiang, Hongmei Xu, Meilin Wang, Zefu Li, Xinyang Su, Xiaoli Li, Zhenzhu Li, Xuexin Han, Effect of infusion speed of 7.5% hypertonic saline on brain edema in patients with craniocerebral injury: An experimental study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Gene(2017), doi:10.1016/j.gene.2018.05.005

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Date: 11th January, 2018.

Dear Editors:

We would like to submit the enclosed manuscript entitled "Effect of infusion speed of 7.5%

hypertonic saline on brain edema in patients with craniocerebral injury: An experimental study", which we wish to be considered for publication in "Gene". We hope this manuscript is suitable for

the article type of "Research paper".

We believe that three aspects of this manuscript will make it interesting to general readers of

"Gene".

Firstly, in recent years, a fact that we cannot hide or deny is that more patients of brain injury

who should have been injected 20% mannitol reduces intracranial pressure. However, Mannitol

has many complications and adverse reactions. Secondly, Now many clinical workers choose 7.5%

hypertonic saline to reduce intracranial pressure and brain edema. More importantly, we hope that

more doctors and scholars would be concerned about that effect of the infusion rate of 7.5%

hypertonic saline on the therapeutic effect. So this randomized controlled trial was designed.

The manuscript hasn't been submitted or published elsewhere. No conflict of interest exits in

this manuscript, and the submission of this manuscript is approved by all the authors. We deeply

appreciate your consideration of our manuscript, and we look forward to receiving comments from

the reviewers. If you have any queries, please don't hesitate to contact me at the address below.

Thank you and best regards.

Yours sincerely,

Affiliation: Affiliated Hospital of Binzhou Medical University, Binzhou, China

Correspondence to: Ms Hongmei XU Affiliated Hospital of Binzhou Medical University, Binzhou,

Shandong 256603, China

E-mail: jzz9008@sina.com

Telephone: 0086-18354359827

## Download English Version:

## https://daneshyari.com/en/article/8644931

Download Persian Version:

https://daneshyari.com/article/8644931

<u>Daneshyari.com</u>