

Accepted Manuscript

Association between FTO gene polymorphisms and type 2 diabetes mellitus, serum levels of apelin and androgen hormones among Iranian obese women

Farzaneh Ghafarian Alipour, Shayan Ziaee, Mohamad Reza Ashoori, Mir Saeid Zakeri, Mohammad Ali Boroumand, Naser Aghamohammadzadeh, Maryam Abbasi-Majdi, Fatemeh Shool, Navid Sarakhs Asbaghi, Abolghasem Mohammadi, Nosratollah Zarghami



PII: S0378-1119(17)30928-9
DOI: doi:[10.1016/j.gene.2017.10.082](https://doi.org/10.1016/j.gene.2017.10.082)
Reference: GENE 42303

To appear in: *Gene*

Received date: 27 March 2017
Revised date: 15 October 2017
Accepted date: 30 October 2017

Please cite this article as: Farzaneh Ghafarian Alipour, Shayan Ziaee, Mohamad Reza Ashoori, Mir Saeid Zakeri, Mohammad Ali Boroumand, Naser Aghamohammadzadeh, Maryam Abbasi-Majdi, Fatemeh Shool, Navid Sarakhs Asbaghi, Abolghasem Mohammadi, Nosratollah Zarghami , Association between FTO gene polymorphisms and type 2 diabetes mellitus, serum levels of apelin and androgen hormones among Iranian obese women. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Gene*(2017), doi:[10.1016/j.gene.2017.10.082](https://doi.org/10.1016/j.gene.2017.10.082)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Association between *FTO* gene polymorphisms and type 2 diabetes mellitus, serum levels of apelin and androgen hormones among Iranian obese women

Farzaneh Ghafarian alipour^{1,2}, Shayan Ziaee³, Mohamad Reza Ashoori^{1,2}, Mir Saeid Zakeri⁴, Mohammad Ali Boroumand³, Naser Aghamohammadzadeh⁵, Maryam Abbasi-Majdi², Fatemeh Shool², Navid Sarakhs Asbaghi⁶, Abolghasem Mohammadi⁷, Nosratollah Zarghami^{1,2*}

¹Nutrition Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Clinical Biochemistry and Laboratory Sciences, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Molecular Pathology, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran.

⁴Department of Neurosurgery, Tabriz University of Medical Sciences, Tabriz, Iran

⁵Department of Endocrinology, Tabriz University of Medical Sciences, Tabriz, Iran

⁶Pardis Clinical Diagnostic Laboratory, University Clinics, Tabriz University of Medical Sciences, Tabriz, Iran

⁷Department of Plant Breeding & Biotechnology, Faculty of Agriculture, University of Tabriz, Iran

*Corresponding author

Prof. Nosratollah Zarghami, Department of Clinical Biochemistry and Laboratory Sciences, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

E-mail: zarghami@tbzmed.ac.ir

Telefax: +98-4133364666

Download English Version:

<https://daneshyari.com/en/article/8646048>

Download Persian Version:

<https://daneshyari.com/article/8646048>

[Daneshyari.com](https://daneshyari.com)