Accepted Manuscript

An HLA class II locus, previously identified by a genome-wide association study, is also associated with susceptibility to pulmonary tuberculosis in a Chinese population

Ruifen Miao, Shuran Huang, Chen Li, Songning Ding, Rong Wang, Ke Xu, Chen Yang, Fei Xu, Haibo Ge

PII: S1567-1348(18)30422-2

DOI: doi:10.1016/j.meegid.2018.06.022

Reference: MEEGID 3561

To appear in: Infection, Genetics and Evolution

Received date: 22 February 2018
Revised date: 22 May 2018
Accepted date: 18 June 2018



Please cite this article as: Ruifen Miao, Shuran Huang, Chen Li, Songning Ding, Rong Wang, Ke Xu, Chen Yang, Fei Xu, Haibo Ge, An HLA class II locus, previously identified by a genome-wide association study, is also associated with susceptibility to pulmonary tuberculosis in a Chinese population. Meegid (2018), doi:10.1016/j.meegid.2018.06.022

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

An HLA class II locus, previously identified by a genome-wide association study, is also associated with susceptibility to pulmonary tuberculosis in a Chinese population

Ruifen Miao ^{1,†,*}, Shuran Huang ^{2,†}, Chen Li ¹, Songning Ding ¹, Rong Wang ¹, Ke Xu ¹, Chen Yang ¹, Fei Xu ¹, Haibo Ge ^{2,*}

¹Department of Chronic Communicable Disease, Nanjing Municipal Center for Disease Control and Prevention, Nanjing 210003, China;

*Corresponding Author: Ruifen Miao, Department of Chronic Communicable Disease, Nanjing Municipal Center for Disease Control and Prevention, 2 Zi-Zhu-Lin, Nanjing 210003, China; Tel (Fax): 86-25-8353-8345; Email: ruifenmiao@126.com; or Haibo Ge, Nanjing Hospital of Chinese Medicine, 1 Jinling Rd., Nanjing 210001, China; Tel (Fax): 86-25-6878-1521; Email:1914326552@qq.com.

Funding: This study was supported by the Nanjing Municipal Program for the Development of Medical Science and Technology (YKK16215).

²Nanjing Hospital of Chinese Medicine, Nanjing 210001, China

[†] These two authors contributed equally to this work

Download English Version:

https://daneshyari.com/en/article/8646590

Download Persian Version:

https://daneshyari.com/article/8646590

Daneshyari.com