

Identifying optic neuritis and transverse myelitis using administrative data

Ruth Ann Marrie MDPH , Okechukwu Ekuma MSc ,
José M.A. Wijnands PhD , Elaine Kingwell PhD , Feng Zhu MSc ,
Yinshan Zhao PhD , John D. Fisk PhD , Charity Evans PhD ,
Helen Tremlett PhD

PII: S2211-0348(18)30290-6
DOI: <https://doi.org/10.1016/j.msard.2018.08.013>
Reference: MSARD 936

To appear in: *Multiple Sclerosis and Related Disorders*

Received date: 16 May 2018

Accepted date: 11 August 2018

Please cite this article as: Ruth Ann Marrie MDPH , Okechukwu Ekuma MSc , José M.A. Wijnands PhD , Elaine Kingwell PhD , Feng Zhu MSc , Yinshan Zhao PhD , John D. Fisk PhD , Charity Evans PhD , Helen Tremlett PhD , Identifying optic neuritis and transverse myelitis using administrative data, *Multiple Sclerosis and Related Disorders* (2018), doi: <https://doi.org/10.1016/j.msard.2018.08.013>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Administrative data can be used to identify individuals with incident optic neuritis
- Administrative data can be used to identify individuals with incident transverse myelitis
- Administrative data can distinguish those with monophasic syndromes from those with an incident presentation of MS
- The average annual incidence of optic neuritis in Manitoba in 2011-2013 was 75.9 per 100,000

Download English Version:

<https://daneshyari.com/en/article/8647254>

Download Persian Version:

<https://daneshyari.com/article/8647254>

[Daneshyari.com](https://daneshyari.com)