


# Environmental Mycobacterium avium subsp. hominissuis have a higher probability to act as a recipient in conjugation than clinical strains

Salma A. Shoulah, Anna M. Oschmann, Abdelfattah Selim, Torsten Semmler, Carsten Schwarz, Elisabeth Kamal, Faysal Hamouda, Elsayed Galila, Wilbert Bitter, Astrid Lewin

| | |
|---|---|
| PII: | S0147-619X(17)30133-6 |
| DOI: | doi:10.1016/j.plasmid.2018.01.003 |
| Reference: | YPLAS 2360 |
| To appear in: | *Plasmid* |
| Received date: | 3 November 2017 |
| Revised date: | 12 January 2018 |
| Accepted date: | 13 January 2018 |

Please cite this article as: Salma A. Shoulah, Anna M. Oschmann, Abdelfattah Selim, Torsten Semmler, Carsten Schwarz, Elisabeth Kamal, Faysal Hamouda, Elsayed Galila, Wilbert Bitter, Astrid Lewin , Environmental Mycobacterium avium subsp. hominissuis have a higher probability to act as a recipient in conjugation than clinical strains. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Yplas(2018), doi:10.1016/j.plasmid.2018.01.003

# Environmental *Mycobacterium avium subsp. hominissuis* have a higher probability to act as a recipient in conjugation than clinical strains

Salma A. Shoulah[1, 2], Anna M. Oschmann[1], Abdelfattah Selim[2], Torsten Semmler[3], Carsten Schwarz[4], Elisabeth Kamal[1], Faysal Hamouda[2], Elsayed Galila[2], Wilbert Bitter[4], Astrid Lewin[1]

[1] Division 16, Mycotic and parasitic Agents and Mycobacteria, Robert Koch Institute, Berlin, Germany

[2]Department of Animal Medicine (Infectious Diseases), Faculty of Veterinary Medicine, Benha University, Egypt

[3]Junior Research Group Microbial Genomics, Robert Koch Institute, Berlin, Germany

[4]Division of Cystic fibrosis/Christiane Herzog Zentrum, Charité-Universitätsmedizin Berlin, Germany

[5]Molecular and Medical Microbiology, VU University & VU University Medical Center, Amsterdam, The Netherlands