Accepted Manuscript

The utility of magnetic resonance imaging for bone involvement in Gaucher disease. Assessing more than bone crises

PhD, Esther Valero MD, Mercedes Roca MD PhD, Pilar Giraldo MD PhD


PII: S1079-9796(16)30219-4

DOI: doi:10.1016/j.bcmd.2016.10.004


Reference: YBCMD 2086

To appear in: Blood Cells, Molecules, and Diseases

Received date: 1 October 2016 Accepted date: 19 October 2016



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

The utility of magnetic resonance imaging for bone involvement in Gaucher Disease. Assessing More than Bone Crises

Marcio Andrade-Campos, MD. PhD.^{1,2} Esther Valero MD³

Mercedes Roca, MD PhD⁴ Pilar Giraldo, MD PhD^{1,2,5}

On behalf of the Spanish group on Gaucher Disease.

- 1- Translational Research Unit, Aragon Health Research Institute (IIS-A), Zaragoza, Spain
- 2- CIBER de Enfermedades Raras, CIBERER, Zaragoza, Spain
- 3- Department of Rheumatology, Lozano Blesa University Hospital, Zaragoza,
- 4- Roca Radiodiagnostic Center, Zaragoza, Spain
- 5- Fundación Española para el Estudio y Terapéutica de la Enfermedad de Gaucher (FEETEG), Zaragoza, Spain

Abstract

Bone effects are the most frequent cause of disability in Gaucher disease (GD). Magnetic resonance imaging (MRI) has improved the study of bone involvement making it possible to measure the extent of infiltration and to identify localized

Download English Version:

https://daneshyari.com/en/article/8648098

Download Persian Version:

https://daneshyari.com/article/8648098

<u>Daneshyari.com</u>