


Accepted Manuscript

Title: Functional characterisation of the actin-depolymerising factor from the apicomplexan *Neospora caninum* (NcADF)

Authors: Luciana Baroni, Luiz M. Pereira, Sutherland K. Maciver, Ana P. Yatsuda



PII: S0166-6851(18)30117-8
DOI: <https://doi.org/10.1016/j.molbiopara.2018.07.008>
Reference: MOLBIO 11141

To appear in: *Molecular & Biochemical Parasitology*

Received date: 15-5-2018
Revised date: 15-7-2018
Accepted date: 15-7-2018

Please cite this article as: Baroni L, Pereira LM, Maciver SK, Yatsuda AP, Functional characterisation of the actin-depolymerising factor from the apicomplexan *Neospora caninum* (NcADF), *Molecular and Biochemical Parasitology* (2018), <https://doi.org/10.1016/j.molbiopara.2018.07.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Functional characterisation of the actin-depolymerising factor from the apicomplexan *Neospora caninum* (NcADF)

Luciana Baroni,^a Luiz M. Pereira,^a Sutherland K. Maciver,^{b,†} Ana P. Yatsuda^{a,†,*}

^a Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Av. do Café, s/n, 14040-930, Ribeirão Preto, SP, Brazil.

^b Centre for Discovery Brain Sciences, Biomedical Sciences, Edinburgh Medical School, University of Edinburgh, Hugh Robson Building, George Square, Edinburgh EH8 9XD, Scotland, United Kingdom.

[†]SKM and APY are co-senior authors.

Corresponding author: Departamento de Análises Clínicas, Bromatológicas e Toxicológicas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto-SP, 14040-930, Brazil. Email address: ayatsuda@fcfrp.usp.br. Tel: +55 16 33154728 (APY).

Highlights

- NcADF affinity for F-actin but not disassembly activity is regulated by pH;
- Higher concentrations of NcADF promote the overshoot effect on actin polymerisation;

Download English Version:

<https://daneshyari.com/en/article/8648323>

Download Persian Version:

<https://daneshyari.com/article/8648323>

[Daneshyari.com](https://daneshyari.com)