

Accepted Manuscript


A segmental duplication in the common ancestor of Brassicaceae is responsible for the origin of the paralogs *KCS6-KCS5*, which are not shared with other angiosperms

Swati Singh, Sandip Das, R. Geeta

PII: S1055-7903(18)30011-3
DOI: <https://doi.org/10.1016/j.ympev.2018.04.018>
Reference: YMPEV 6128

To appear in: *Molecular Phylogenetics and Evolution*

Received Date: 10 January 2018
Revised Date: 11 April 2018
Accepted Date: 11 April 2018



Please cite this article as: Singh, S., Das, S., Geeta, R., A segmental duplication in the common ancestor of Brassicaceae is responsible for the origin of the paralogs *KCS6-KCS5*, which are not shared with other angiosperms, *Molecular Phylogenetics and Evolution* (2018), doi: <https://doi.org/10.1016/j.ympev.2018.04.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A segmental duplication in the common ancestor of Brassicaceae is responsible for the origin of the paralogs *KCS6-KCS5*, which are not shared with other angiosperms

Swati Singh, Sandip Das, R. Geeta*

Department of Botany, University of Delhi, Delhi 110007.

*Corresponding author - rgeeta@botany.du.ac.in

Key words: *KCS5*, *KCS6*, Brassicaceae, Cuticular wax

Download English Version:

<https://daneshyari.com/en/article/8648853>

Download Persian Version:

<https://daneshyari.com/article/8648853>

[Daneshyari.com](https://daneshyari.com)