

Accepted Manuscript

Systemic monocyte chemotactic protein-1 inhibition modifies renal macrophages and restores glomerular endothelial glycocalyx and barrier function in diabetic nephropathy

Margien G.S. Boels, Angela Koudijs, M. Cristina Avramut, Wendy M.P.J. Sol, Gangqi Wang, Annemarie M. van Oeveren-Rietdijk, Anton Jan van Zonneveld, Hetty C. de Boer, Johan van der Vlag, Cees van Kooten, Dirk Eulberg, Bernard M. van den Berg, Daphne H.T. IJpelaar, Ton J. Rabelink

PII: S0002-9440(17)30006-8

DOI: [10.1016/j.ajpath.2017.07.020](https://doi.org/10.1016/j.ajpath.2017.07.020)

Reference: AJPA 2706

To appear in: *The American Journal of Pathology*


Received Date: 0002-9440 February 0002-9440

Revised Date: 0002-9440 February 0002-9440

Accepted Date: 5 July 2017

Please cite this article as: Boels MGS, Koudijs A, Avramut MC, Sol WMPJ, Wang G, van Oeveren-Rietdijk AM, van Zonneveld AJ, de Boer HC, van der Vlag J, van Kooten C, Eulberg D, van den Berg BM, IJpelaar DHT, Rabelink TJ, Systemic monocyte chemotactic protein-1 inhibition modifies renal macrophages and restores glomerular endothelial glycocalyx and barrier function in diabetic nephropathy, *The American Journal of Pathology* (2017), doi: 10.1016/j.ajpath.2017.07.020.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Systemic monocyte chemotactic protein-1 inhibition modifies renal macrophages and restores glomerular endothelial glycocalyx and barrier function in diabetic nephropathy

Running head: MCP-1 inhibition restores glycocalyx

Margien G.S. Boels¹, Angela Koudijs¹, M. Cristina Avramut², Wendy M.P.J. Sol¹, Gangqi Wang¹, Annemarie M. van Oeveren-Rietdijk¹, Anton Jan van Zonneveld¹, Hetty C. de Boer¹, Johan van der Vlag³, Cees van Kooten¹, Dirk Eulberg⁴, Bernard M. van den Berg¹, Daphne H.T. Ijpelaar¹ Ton J. Rabelink¹

¹ Eindhoven Laboratory for Vascular and Regenerative Medicine, Department of Internal Medicine, Division of Nephrology, Leiden University Medical Center, Leiden, The Netherlands

² Department of Molecular Cell Biology, Leiden University Medical Center, Leiden, The Netherlands

³ Department of Nephrology, Radboud Institute for Molecular Life Sciences, Radboud University Medical Center, Nijmegen, The Netherlands

⁴ NOXXON Pharma AG, Berlin, Germany

Corresponding author:

Ton J. Rabelink

Leiden University Medical Center, Internal Medicine, Department of Nephrology
Albinusdreef 2, 2333 ZA, Leiden, The Netherlands

Fax: +3171 526 6868, Phone: +3171 526 2148, e-mail: a.j.rabelink@lumc.nl

Number of text pages: 17

Number of figures: 5

Funding: Supported by the Dutch Kidney Foundation: Kolff Grant 14OKG06 (D.I.) and Glycoren consortium (CP09.03; T.R.).

Disclosures: D.E. is employed at NOXXON Pharma AG.

Download English Version:

<https://daneshyari.com/en/article/8652130>

Download Persian Version:

<https://daneshyari.com/article/8652130>

[Daneshyari.com](https://daneshyari.com)