



## CASE REPORT

# Retrograde catheterization of the SMA on a B-EVAR – Case report



Gonçalo Queiroz de Sousa<sup>a,\*</sup>, Ruy Fernandes e Fernandes<sup>a</sup>, Luís Mendes Pedro<sup>a</sup>, Pedro Garrido<sup>a</sup>, Luís Silvestre<sup>a</sup>, Eric Verhoeven<sup>b</sup>, José Fernandes e Fernandes<sup>a</sup>

<sup>a</sup> *Clínica Universitária de Cirurgia Vascular, Hospital de Santa Maria – Centro Hospitalar Lisboa Norte, Faculdade de Medicina da Universidade de Lisboa, Centro Académico de Medicina de Lisboa, Lisboa, Portugal*

<sup>b</sup> *Department of Vascular and Endovascular Surgery, Paracelsus Medical University, Nuremberg, Germany*

Received 7 June 2016; accepted 9 September 2016

Available online 28 October 2016

### KEYWORDS

Retrograde catheterization;  
Superior mesenteric artery;  
Fenestrated evar;  
Branched evar;  
Thoracoabdominal aneurysm

**Abstract** Endovascular treatment of complex thoracoabdominal aneurysms with branched and fenestrated grafts (B/F-EVAR) has become the first option for patients with suitable anatomy and very high risk for open surgery, who would likely be refused for open repair. We present a case of a patient with a type III thoracoabdominal aneurysm submitted to endovascular repair with a custom-made endograft with branches to the celiac trunk and SMA, and a fenestration to the left renal artery. During the procedure there was the need to do a laparotomy to allow for retrograde catheterization of the SMA, with technical success. In this case, the retrograde access was of utmost importance for the completion of the procedure. Different techniques for target vessel catheterization, such as the one we describe, should be part of a vascular surgeon's resources in B/F-EVAR procedures.

© 2016 Sociedade Portuguesa de Angiologia e Cirurgia Vascular. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### PALAVRAS CHAVE

Cateterização retrógrada;  
Artéria mesentérica superior;  
Endoprótese fenestrada;

### Cateterização retrógrada da AMS como recurso em B-EVAR – Caso clínico

**Resumo** O tratamento endovascular de aneurismas toraco-abdominais complexos com endopróteses ramificadas e/ou fenestradas (B/F-EVAR), coloca-se como primeira opção em doentes com risco cirúrgico muito elevado e anatomia favorável, e que potencialmente seriam recusados para tratamento cirúrgico convencional. Os autores apresentam um caso clínico de um doente com um aneurisma da aorta toraco-abdominal tipo III submetido a B/F-EVAR com endoprótese *custom-made* com ramificações para o tronco celiaco e artéria mesentérica superior, e fenestração para a artéria renal esquerda. Durante o procedimento houve necessidade de

\* Corresponding author.

E-mail address: [queiroz.sousa@gmail.com](mailto:queiroz.sousa@gmail.com) (G. Queiroz de Sousa).

Endoprótese  
ramificada;  
Aneurisma  
toraco-abdominal

realizar laparotomia mediana para cateterização retrógrada da AMS, concretizada com sucesso. Neste caso, a cateterização retrógrada da AMS foi fulcral para o sucesso técnico da intervenção. Técnicas de cateterização alternativas como a descrita devem ser parte do armamentário de recurso na execução de casos de B/F-EVAR.

© 2016 Sociedade Portuguesa de Angiologia e Cirurgia Vascular. Publicado por Elsevier España, S.L.U. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Since endovascular aneurysm repair (EVAR) was introduced,<sup>1</sup> technology has evolved and custom-made fenestrated grafts have been made available since 1999,<sup>2</sup> which allowed to take the endovascular repair to even more complex aortic aneurysms, namely juxtarenal, suprarenal and thoracoabdominal (TAAA) ones. Besides fenestrations, custom-made branched grafts have also been developed and used to repair TAAA.<sup>3</sup> This has become a feasible technique with promising short and mid-term results reported in high volume centers.<sup>4,5</sup> These techniques bring a possibility of endovascular repair to patients with aneurysms of great anatomical complexity whose open surgical treatment would probably require a supra-celiac clamping of the aorta.

However, despite careful planning of the procedures, difficulties in target vessel catheterization may occur during branched and fenestrated EVAR (B/F-EVAR), and bailout techniques can be used.<sup>6,7</sup> We report a case of a successful retrograde catheterization of the superior mesenteric artery (SMA) using supra-umbilical laparotomy during B/F-EVAR.

## Clinical case

A 70-year-old caucasian male, with known history of chronic obstructive pulmonary disease (COPD) with severe bronchic and bronchiolar obstruction on respiratory tests, stage 3<sup>8</sup> chronic kidney disease with an eGFR of 38.5 mL/min/1.73 m<sup>2</sup>, arterial hypertension and smoking in the past, was sent from another Hospital's outpatient clinic to our Vascular Surgery Department with the diagnosis of type III TAAA with 60 mm diameter (Fig. 1A). The patient was considered of high risk for open surgery, and proposed for endovascular repair with a custom-made graft.

Regarding planning for B/F-EVAR, several technical details had to be considered due to the patient's anatomy, namely: the tight double angulation of the aorta below the origin of the celiac trunk (CT) and SMA (Fig. 1B); a thin right renal artery (RRA) with an atrophic kidney (Fig. 2A); two left renal arteries (LRA), with the thinner one going to lower kidney pole (Fig. 2B and C); and the fairly low diameter of the aorta at the level of the renal arteries.

After thorough planning, a customized Cook Zenith® graft (William A. Cook Australia Ltd., Brisbane, Australia) was designed, and included a proximal tapered thoracic component, a main body with branches to the CT (8 × 18 mm, diameter × length) and SMA (8 × 18 mm) and a fenestration (6 × 8 mm, width × height) for the main LRA, and a distal bifurcated graft plus contralateral limb. As part

of the plan for this main procedure, the patient underwent coil embolization of the RRA and thinner LRA 2 weeks before B/F-EVAR, without any complications. It was not feasible to build a graft with 2 fenestrations for the left renal arteries because they arose from the aorta at almost the same level and there was only 1 h 45 min between them, and a minimum of 2 h 15 min was needed for this to be possible.

The patient underwent the procedure under general anesthesia, a spinal catheter for cerebrospinal fluid (CSF) drainage was placed the day before, contrast-induced kidney injury prophylaxis was done with pre-procedural intravenous hydration with saline and N-acetylcysteine, and both femoral and left axillary arteries were used as access using surgical exposure. The main LRA and CT were catheterized without issues. We were not able to catheterize the SMA using either the axillary or the femoral access, mainly due to the angulation of the aorta which led practically to the occlusion of the SMA's ostium by the main graft's body. After all the failed attempts to catheterize the SMA endovascularly, a median laparotomy was performed, and the SMA was retrogradely catheterized. The guidewire easily entered the endograft's branch retrogradely, and the SMA was stented with success (Fig. 3A–D) which allowed for completion of the B/F-EVAR, with an estimated partial bowel ischemia time of 1 h.

The patient was transferred to the intensive care unit (ICU), where he stayed for 6 days. There was a transient deterioration of kidney function, AKIN II, that recovered to pre-operative values with medical therapy alone and no need for dialysis. In the first 8 h after the surgery, the patient had transient intra-abdominal hypertension (IAH) with a pressure of 24 mmHg and high lactates of 40 mg/dL, that resolved with medical treatment, namely sedation and analgesia. 24 h after the surgery the patient developed paraparesis which required vasoactive support with norepinephrine in order to achieve mean arterial pressures (MAP) of 90–100 mmHg, and lowering the spinal cord drain pressure to 5 mmHg, to completely revert the neurological deficits successfully within another 24 h; a neurological evaluation was done hourly in the first 12 h. The patient was discharged 14 days after surgery. The first control CT angiogram was only at 6 months considering the patient's chronic kidney disease, and showed both branches and LRA stent patent (Fig. 4A and B).

## Discussion

Fenestrated and branched stent grafts are becoming more frequently used in our department, which allows us to offer

Download English Version:

<https://daneshyari.com/en/article/8652218>

Download Persian Version:

<https://daneshyari.com/article/8652218>

[Daneshyari.com](https://daneshyari.com)