


Accepted Manuscript

Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally

Milton Fabian Suárez-Ortegón, Stela McLachlan, Anna H. Price, Mercé Fernández-Balsells, Josep Franch-Nadal, Manel Mata-Cases, Joan Barrot-de la Puente, Xavier Mundet-Tudurí, Didac Mauricio, Wifredo Ricart, Sarah H. Wild, Mark W.J. Strachan, Jackie F. Price, José-Manuel Fernández-Real



PII: S0021-9150(18)30153-9

DOI: [10.1016/j.atherosclerosis.2018.03.028](https://doi.org/10.1016/j.atherosclerosis.2018.03.028)

Reference: ATH 15436

To appear in: *Atherosclerosis*

Received Date: 7 December 2017

Revised Date: 27 February 2018

Accepted Date: 15 March 2018

Please cite this article as: Suárez-Ortegón MF, McLachlan S, Price AH, Fernández-Balsells Mercé, Franch-Nadal J, Mata-Cases M, Barrot-de la Puente J, Mundet-Tudurí X, Mauricio D, Ricart W, Wild SH, Strachan MWJ, Price JF, Fernández-Real José-Manuel, Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally, *Atherosclerosis* (2018), doi: [10.1016/j.atherosclerosis.2018.03.028](https://doi.org/10.1016/j.atherosclerosis.2018.03.028).

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally

Milton Fabian Suárez-Ortegón ^{1,2*}, Stela McLachlan ¹, Anna H. Price ¹, Mercé Fernández-Balsells³, Josep Franch-Nadal⁴⁻⁶, Manel Mata-Cases^{4,6,7}, Joan Barrot-de la Puente⁴, Xavier Mundet-Tuduri⁴, Didac Mauricio⁶, Wifredo Ricart³, Sarah H. Wild ¹, Mark W. J. Strachan ⁸, Jackie F. Price ¹, José-Manuel Fernández-Real ^{3*}

1. Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, Scotland-UK
2. Nutrition Group, Universidad del Valle, Cali-Colombia
3. Department of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de Girona (IdIBGi), CIBEROBN (CB06/03/010) and Instituto de Salud Carlos III (ISCIII), Girona- Spain
4. DAP-Catgroup. Unitat de Suport a la Recerca Barcelona Ciutat, Institut Universitari d'Investigació en Atenció Primària Jordi Gol (IDIAP Jordi Gol), Barcelona-Spain
- 5 Primary Health Care Center Raval Sud, Gerència d'Àmbit d'Atenció Primària Barcelona Ciutat, Institut Català de la Salut, Barcelona-Spain
- 6-Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas (CIBERDEM), Instituto de Salud Carlos III (ISCIII)-Spain
- 7-Primary Health Care La Mina (Sant Adrià de Besò). Gerència d'Àmbit d'Atenció Primària Barcelona Ciutat, Institut Català de la Salut, Barcelona-Spain
- 8- Metabolic Unit, Western General Hospital, Edinburgh, Scotland-UK

*Corresponding authors:

MF Suarez Ortegón, Centre for Population Health Sciences. College of Medicine and Veterinary Medicine. University of Edinburgh. Teviot Place. Edinburgh EH8 9AG. United Kingdom. Email: Milton.Suarez@ed.ac.uk.

JM Fernández-Real, Department of Diabetes, Endocrinology and Nutrition, Hospital of Girona "Dr Josep Trueta", Carretera de França s/n, 17007, Girona, Spain. Email: jmfreal@idibgi.org.

Download English Version:

<https://daneshyari.com/en/article/8656895>

Download Persian Version:

<https://daneshyari.com/article/8656895>

[Daneshyari.com](https://daneshyari.com)