

H.E.L.P apheresis exerts long term effects on the capacity of circulating proangiogenic cells

Jan Sradnick, Sergey Tselmin, Andrea Wagner, Ulrich Julius, Vladimir Todorov, Christian Hugo, Bernd Hohenstein

PII: S1567-5688(17)30087-9

DOI: 10.1016/j.atherosclerosissup.2017.05.045

Reference: ATHSUP 337

To appear in: *Atherosclerosis (Supplements) (Component)*

Please cite this article as: Sradnick J, Tselmin S, Wagner A, Julius U, Todorov V, Hugo C, Hohenstein B, H.E.L.P apheresis exerts long term effects on the capacity of circulating proangiogenic cells, *Atherosclerosis (Supplements) (Component)* (2017), doi: 10.1016/j.atherosclerosissup.2017.05.045.

H.E.L.P apheresis exerts long term effects on the capacity of circulating proangiogenic cells

Jan Sradnick[1], Sergey Tselmin[2], Andrea Wagner[1], Ulrich Julius[2], Vladimir Todorov[1], Christian Hugo[1] and Bernd Hohenstein [1,2]

[1]Division of Nephrology and , [2] Extracorporeal Treatment and Apheresis Center, Department of Internal Medicine III, University Hospital and Faculty of Medicine Carl Gustav Carus at the Technische Universität, Dresden, Germany

Corresponding author

Prof. Dr. Bernd Hohenstein

Division of Nephrology and Extracorporeal Treatment and Apheresis Center

Department of Internal Medicine III

University Hospital and Faculty of Medicine Carl Gustav Carus

at the Technische Universität Dresden

Fetscherstraße 74

01307 Dresden, Germany

Phone: +49 351 458 3075

Fax: +49 351 458 5333

bernd.hohenstein@ukdd.de

Keywords: lipoprotein apheresis, PAC, EPC, LDL-C, lipoprotein(a), endothelial repair