

CASE REPORT

Restenosis of Aorto-renal Venous Grafts: Report of Two Patients Treated by Endovascular Stenting

Miguel Lemos Gomes^{a,b,*}, Alice Lopes^{a,b}, Gonçalo Sobrinho^{a,b,c}, Luis Mendes Pedro^{a,b,c}

^a Vascular Surgery, Heart and Vessels Department, Hospital de Santa Maria (CHLN), Lisbon, Portugal

^b Lisbon Academic Medical Centre, Lisbon, Portugal

^c Medical School, University of Lisbon, Lisbon, Portugal

Introduction: Renal venous graft restenosis is an uncommon event usually associated with significant clinical impact. Its treatment by endovascular stenting is seldom reported in the literature.

Report: Two cases of successful stenting for restenosis in aorto-renal venous grafts are described, detailing the technique and in one case reporting for the first time the use of a covered stent in this condition.

Discussion: Technical success may be achieved with proper material selection for the patient's anatomy and with dilation at relatively high pressures. The use of a covered stent may provide extra safety when treating vein grafts.

© 2018 The Authors. Published by Elsevier Ltd on behalf of European Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Article history: Received 1 March 2018, Revised 30 June 2018, Accepted 2 July 2018,

Keywords: Renal artery disease, Restenosis, Stent graft

INTRODUCTION

Based on Goldblatt's innovative work defining the pathophysiology of renovascular disease in 1934,¹ surgical treatment has evolved from nephrectomy to open surgical procedures, mostly bypass techniques using venous or synthetic grafts.^{2–4} Presently, the preferred technique for renal artery revascularisation is endovascular angioplasty or stenting, a method introduced by Gruntzig in 1978,⁵ and currently used in many patients.

One complication of renal bypass grafts is restenosis, which has been reported as having an incidence of 3.4%.⁶ Restenosis of renal artery bypasses is always difficult to manage and, surprisingly, there are few descriptions of its treatment by an endovascular procedure, with only three cases reported in the literature.^{7,8}

The aim of this study is to report two cases of renal bypass graft stenoses successfully treated by endovascular stenting, and to detail and discuss the technical aspects of the procedure.

CASE 1

In 2006, a 72 year old woman with arterial hypertension underwent an aorto-bifemoral bypass for occlusive aortic disease associated with prosthetic revascularisation of the

superior mesenteric artery and bilateral venous bypass to both renal arteries due to tight ostial stenosis (inflow from the aortic graft). After the procedure, renal function and hypertension normalised. During follow up, bilateral ureteral stenting was needed to manage ureteral compression.

Two years after the initial surgery, renal function deteriorated (blood creatinine increased from 0.7 mg/dL to 2.7 mg/dL) and, to achieve proper blood pressure control, the number of antihypertensive medications prescribed increased from two to four. An aortic angiogram documented severe stenosis of both venous renal grafts.

Under general anaesthesia and by exposure of the right superficial femoral artery to avoid exposure of the Dacron limb (thus reducing the chance of an infectious complication), a 7 Fr sheath was inserted and a guiding catheter (Mach1™ Guide Catheter; Boston Scientific, Natick, MA, USA) was advanced to the main body of the aortic graft. Catheterisation of the venous conduits was performed with a 0.035 Magic Torque™ Guidewire (Boston Scientific) followed by deployment of balloon expandable stents in the proximal anastomosis on the right and at the middle portion of the bypass on the left (Express™ SD Renal Monorail™ 6 × 18 mm on the right and 5 × 19 mm on the left; Boston Scientific [Fig. 1]). The procedure was uneventful. After 8 years of follow up the patient remains asymptomatic with blood creatinine stable at 1.4 mg/dL, and is currently taking two antihypertensive drugs. A recent computed tomographic angiogram (CTA) showed patency of both stents (Fig. 2).

CASE 2

A 69 year old man with ischaemic heart disease (having undergone a coronary artery bypass graft 13 years

* Corresponding author. Vascular Surgery, Heart and Vessels Department, Hospital de Santa Maria (CHLN), Lisbon, Portugal.

E-mail address: gomes.l.miguel@gmail.com (Miguel Lemos Gomes).

2405-6553/© 2018 The Authors. Published by Elsevier Ltd on behalf of European Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.ejvssr.2018.07.001>




Figure 1. Pre-occlusive stenosis of the right aorto-renal bypass and moderate stenosis of the left aorto-renal bypass; the angiographic result after placement of balloon expandable stents.

previously), hypertension, dyslipidaemia, chronic kidney disease (stage III/V), and peptic ulcer had open surgery for treatment of a juxtarenal aortic aneurysm in 2013. This procedure included aorto-bifemoral graft interposition, associated with right renal artery bypass with the great saphenous vein (inflow from the aortic graft), owing to a severe stenosis in the origin of that artery (atrophic left

kidney). The recovery was uneventful, with improvement of hypertension and stabilisation of renal function (blood creatinine level 1.4mg/dL [stage III/V]).

After three years of follow up, a deterioration in renal function was observed (blood creatinine increased from 1.4 mg/dL to 2.5mg/dL) and a CTA revealed severe stenosis of the renal venous bypass ([Fig. 3](#)). Owing to the recent




Figure 2. Computed tomography angiography showing patency of both stents.

Download English Version:

<https://daneshyari.com/en/article/8659187>

Download Persian Version:

<https://daneshyari.com/article/8659187>

[Daneshyari.com](https://daneshyari.com)